arXiv:1303.0827v1 [math.DG] 4 Mar 2013

CRITICAL METRICS ON CONNECTED SUMS OF EINSTEIN FOUR-MANIFOLDS

MATTHEW J. GURSKY AND JEFF A. VIACLOVSKY

Abstract. We develop a gluing procedure designed to obtain canonical metrics on connected sums of Einstein four-manifolds. The main application is an existence result, using two well-known Einstein manifolds as building blocks: the Fubini-Study metric on $\mathbb{CP}^2$ and the product metric on $S^2 \times S^2$. Using these metrics in various gluing configurations, critical metrics are found on connected sums for a specific Riemannian functional, which depends on the global geometry of the factors. Furthermore, using certain quotients of $S^2 \times S^2$ as one of the gluing factors, critical metrics on several non-simply-connected manifolds are also obtained.

Contents

1. Introduction 1
2. The building blocks 10
3. The nonlinear map 17
4. Cokernel on a compact manifold 26
5. Cokernel on an asymptotically flat manifold 29
6. Asymptotics of the cokernel 36
7. Some auxiliary linear equations 40
8. Computation of the leading term 46
9. Naïve approximate metric 55
10. Refined approximate metric 59
11. Lyapunov-Schmidt reduction 68
12. Completion of proofs 78

Appendix A. Proof of Proposition 8.5 85

Appendix B. Non-simply-connected examples 89

References 89

1. Introduction

A Riemannian manifold (M^4, g) in dimension four is critical for the Einstein-Hilbert functional

$$\mathcal{R}(g) = Vol(g)^{-1/2} \int_M R_g dV_g, \tag{1.1}$$

Date: March 4, 2013.

where R_g is the scalar curvature, if and only if it satisfies

$$Ric(g) = \lambda \cdot g, \tag{1.2}$$

where λ is a constant; such Riemannian manifolds are called *Einstein manifolds.* Non-collapsing limits of Einstein manifolds have been studied in great depth [And89, BKN89, Tia90]. In particular, with certain geometric conditions, the limit space is an orbifold, with asymptotically locally Euclidean (ALE) spaces bubbling off at the singular points. A natural question is whether it is possible to reverse this process: can one start with the limit space, and glue on a bubble in order to obtain an Einstein metric? A recent article of Olivier Biquard makes great strides in the Poincaré-Einstein setting [Biq11]. In this work it is shown that a $\mathbb{Z}/2\mathbb{Z}$-orbifold singularity p of a non-degenerate Poincaré-Einstein orbifold (M, g) has a Poinaré-Einstein resolution obtained by gluing on an Eguchi-Hanson metric if and only if the condition

$$\det(\mathbf{R}^+(p)) = 0 \tag{1.3}$$

is satisfied, where $\mathbf{R}^+(p) : \Lambda^2_+ \to \Lambda^2_+$ is the purely self-dual part of the curvature operator at p. The self-adjointness of this gluing problem is overcome by the freedom of changing the boundary data of the Poincaré-Einstein metric.

However, there is not much known about gluing compact manifolds together in the Einstein case. In this work, we will replace the Einstein equations with a generalization of the Einstein condition. Namely, we ask whether it is possible to glue together Einstein metrics and produce a critical point of a certain Riemannian functional generalizing the Einstein-Hilbert functional. It turns out that there is a family of such functionals; this gives an extra parameter which will allow us to overcome the self-adjointness of this problem. The particular functional will then depend on the global geometry of the gluing factors.

To describe the functionals, let M be a closed manifold of dimension 4. We will consider functionals on the space of Riemannian metrics $\mathcal{M}$ which are quadratic in the curvature. Such functionals have also been widely studied in physics under the name "fourth-order," "critical," or "quadratic" gravity; see for example [LP11, Mal11, Sch07, Ste78]. In previous work, the authors have studied rigidity and stability properties of Einstein metrics for quadratic curvature functionals [GV11]; these results will play a crucial rôle in this paper.

Using the standard decomposition of the curvature tensor Rm into the Weyl, Ricci and scalar curvature curvature components (denoted by W, Ric, and R, respectively), a basis for the space of quadratic curvature functionals is

$$\mathcal{W} = \int |W|^2 \, dV, \quad \rho = \int |Ric|^2 \, dV, \quad \mathcal{S} = \int R^2 \, dV, \tag{1.4}$$

where we use the tensor norm. In dimension four, the Chern-Gauss-Bonnet formula

$$32\pi^2 \chi(M) = \int |W|^2 \, dV - 2 \int |Ric|^2 \, dV + \frac{2}{3} \int R^2 \, dV \tag{1.5}$$

implies that ρ can be written as a linear combination of the other two (plus a topological term). Consequently, we will be interested in the functional

$$\mathcal{B}_t[g] = \int |W|^2 \, dV + t \int R^2 \, dV \tag{1.6}$$

(with $t = \infty$ formally corresponding to $\int R^2 dV$).

The Euler-Lagrange equations of $\mathcal{B}_t$ are given by

$$B^t \equiv B + tC = 0, \tag{1.7}$$

where B is the *Bach tensor* defined by

$$B_{ij} \equiv -4\Big(\nabla^k \nabla^l W_{ikjl} + \frac{1}{2} R^{kl} W_{ikjl}\Big) = 0, \tag{1.8}$$

and C is the tensor defined by

$$C_{ij} = 2\nabla_i \nabla_j R - 2(\Delta R) g_{ij} - 2R R_{ij} + \frac{1}{2} R^2 g_{ij}. \tag{1.9}$$

It follows that any Einstein metric is critical for $\mathcal{B}_t$ [Bes87]. We will refer to such a critical metric as a *B^t-flat metric*. Note that by taking a trace of (1.7), it follows that the scalar curvature of a B^t-flat metric on a compact manifold is necessarily constant. Therefore a B^t-flat metric satisfies the equation

$$B = 2tR \cdot E, \tag{1.10}$$

where E denotes the traceless Ricci tensor. That is, the Bach tensor is a constant multiple of the traceless Ricci tensor.

The convergence results described above for Einstein metrics were generalized to systems of the form

$$\Delta Ric = Rm * Ric \tag{1.11}$$

(of which (1.10) is a special case) in [TV05a, TV05b, TV08]. In particular, with certain geometric conditions, non-collapsing sequences of metrics satisfying an equation of the form (1.11) have orbifold limits. Again, the natural question is whether it is possible to reverse this bubbling process.

The analogous gluing problem for the anti-self-dual equations $W^+ = 0$ in dimension four has been very successful [DF89, Flo91, Tau92, KS01, AV12a]. However, gluing for the B^t-flat equations is much more difficult because, as in the Einstein case, this is a self-adjoint problem. The parameter t is the key to overcoming this difficulty.

We point out that the linearization of the B^t-flat equation (1.7) is not elliptic due to diffeomorphism invariance. It will be necessary to "gauge" the equation in order to work with an elliptic operator. This is analogous to the Bianchi gauge for the Einstein equations. The details of this gauging process appear in Section 3.

The main building blocks in this paper are the Fubini-Study metric $(\mathbb{CP}^2, g_{FS})$, and $(S^2 \times S^2, g_{S^2 \times S^2})$, the product of 2-dimensional spheres with unit Gauss curvature. Both are Einstein, so are B^t-flat for all $t \in \mathbb{R}$. A key result used in this paper is rigidity of these metrics for certain ranges of t, which was proved in our previous work [GV11]. That is, these metrics admit no non-trivial infinitesimal B^t-flat deformations for certain ranges of t (other than scalings). These rigidity properties will be discussed in Section 4.

1.1. Green's function metric.

Recall that the conformal Laplacian is the operator

$$Lu = -6\Delta u + Ru, \tag{1.12}$$

where our convention is to use the analyst's Laplacian (which has negative eigenvalues). If (M, g) is compact and has positive scalar curvature, then for any $p \in M$, there exists a unique positive solution to the equation

$$LG = 0 \quad \text{on } M \setminus \{p\} \tag{1.13}$$

$$G = \rho^{-2}(1 + o(1)) \tag{1.14}$$

as $\rho \to 0$, where ρ is geodesic distance to the basepoint p, which is called the Green's function. Denote $N = M \setminus \{p\}$ with metric $g_N = G^2 g_M$. The metric g_N is scalar-flat and asymptotically flat of order 2. Recall the mass of an AF space is defined by

$$\text{mass}(g_N) = \lim_{R\to\infty} \omega_3^{-1} \int_{S(R)} \sum_{i,j} (\partial_i g_{ij} - \partial_j g_{ii})(\partial_i \lrcorner\, dV), \tag{1.15}$$

with $\omega_3 = Vol(S^3)$.

A crucial point is the following: if (M, g) is Bach-flat, then from conformal invariance of the Bach tensor, (N, g_N) is also Bach-flat. Also, since the Green's function is used as the conformal factor, g_N is scalar-flat. Consequently, g_N is B^t-flat for all $t \in \mathbb{R}$.

The Green's function metric of the Fubini-Study metric $\hat{g}_{FS}$ is also known as the Burns metric, and is completely explicit, with mass given by

$$\text{mass}(\hat{g}_{FS}) = 2. \tag{1.16}$$

However, the Green's function metric $\hat{g}_{S^2\times S^2}$ of the product metric does not seem to have a known explicit description. We will denote

$$m_1 = \text{mass}(\hat{g}_{S^2\times S^2}). \tag{1.17}$$

By the positive mass theorem of Schoen-Yau, $m_1 > 0$ [SY79, SY81]. We note that since $S^2 \times S^2$ is spin, this also follows from Witten's proof of the positive mass theorem [Wit81].

1.2. The gluing procedure.

Let (Z, g_Z) and (Y, g_Y) be Einstein manifolds, and assume that g_Y has positive scalar curvature. Choose basepoints $z_0 \in Z$ and $y_0 \in Y$. Convert (Y, g_Y) into an asymptotically flat (AF) metric (N, g_N) using the Green's function for the conformal Laplacian based at y_0. As pointed out above, g_N is B^t-flat for any t.

Let $a > 0$ be small, and consider $Z \setminus B(z_0, a)$. Scale the compact metric to $(Z, \tilde{g} = a^{-4} g_Z)$. Attach this metric to the metric $(N \setminus B(a^{-1}), g_N)$ using cutoff functions near the boundary, to obtain a smooth metric on the connect sum $Z \# \overline{Y}$. Since both g_Z and g_N are B^t-flat, this metric is an "approximate" B^t-flat metric, with vanishing

B^t tensor away from the "damage zone", where cutoff functions were used. This construction is described in detail in Section 9, and is illustrated in Figure 1.1.



FIGURE 1.1. The approximate metric.

This "naïve" approximate metric is too rough for our purposes – the size of the B^t tensor is an order too large in the damage zone. A refinement of this approximate metric is found by solving linear equations on each piece to make the metrics match up to highest order. The B^t tensor of the refined metric is now an order of magnitude smaller. This step is inspired by the recent work of Biquard in the Einstein case which was mentioned above [Biq11]. These auxiliary linear equations are solved in Section 7, and the refined approximate metric is constructed in Section 10.

Lyapunov-Schmidt reduction is then used to reduce the problem from an infinite-dimensional problem to a finite-dimensional one. That is, the problem of finding a B^t-flat metric is reduced to finding a zero of the Kuranishi map, which is a mapping between finite-dimensional spaces. This reduction is carried out in Section 11.

For the general gluing problem, even if the pieces are rigid, there can be nonzero infinitesimal kernel elements due to the presence of gluing parameters. In general, there are infinitesimal kernel elements corresponding geometrically to freedom of scaling the AF space, rotating the gluing factor, and moving the base points of the gluing. The leading term of the Kuranishi map corresponding to the scaling parameter, denoted by $\lambda_1(a)$, is given by:

Theorem 1.1. *As $a \to 0$, then for any $\epsilon > 0$,*

$$\lambda_1(a) = \Big(\frac{2}{3} W(z_0) \circledast W(y_0) + 4tR(z_0)\mathrm{mass}(g_N)\Big)\omega_3 a^4 + O(a^{6-\epsilon}), \tag{1.18}$$

where $\omega_3 = Vol(S^3)$, and the product of the Weyl tensors is given by

$$W(z_0) \circledast W(y_0) = \sum_{ijkl} W_{ijkl}(z_0)(W_{ijkl}(y_0) + W_{ilkj}(y_0)), \tag{1.19}$$

where $W_{ijkl}(\cdot)$ *denotes the components of the Weyl tensor in a normal coordinate system at the corresponding point.*

We note that the product $\circledast$ depends upon the coordinate systems chosen, and therefore in general depends upon a rotation parameter, and obviously on the base points of the gluing.

1.3. **Simply-connected examples.** In the case either of the factors are $(\mathbb{CP}^2, g_{FS})$ or $(S^2 \times S^2, g_{S^2\times S^2})$, Theorem 1.1 implies an existence theorem. Since these manifolds are toric, we can use the torus action plus a certain discrete symmetry, called a diagonal symmetry, to eliminate all gluing parameters except for the scaling parameter. Theorem 1.1 will then allow us to obtain critical metrics on the following manifolds "near" the indicated approximate metric:

- (i) $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$; the Fubini-Study metric with a Burns metric attached at one fixed point. This case admits a $U(2)$-action.
- (ii) $S^2 \times S^2 \# \overline{\mathbb{CP}}^2 = \mathbb{CP}^2 \# 2\overline{\mathbb{CP}}^2$; the product metric on $S^2 \times S^2$ with a Burns metric attached at one fixed point. Alternatively, we can view this as the Fubini-Study metric on $\mathbb{CP}^2$, with a Green's function $S^2 \times S^2$ metric attached at one fixed point. For this topology, we will therefore construct two different critical metrics.
- (iii) $2\# S^2 \times S^2$; the product metric on $S^2 \times S^2$ with a Green's function $S^2 \times S^2$ metric attached at one fixed point.

More precisely, we have

Theorem 1.2. *In each of the above cases, a* B^t*-flat metric exists for some* t *near the critical value of*

$$t_0 = \frac{-1}{6R(z_0)\mathrm{mass}(g_N)} W(z_0) \circledast W(y_0). \tag{1.20}$$

Furthermore, this metric is invariant under the indicated action(s).

The proof of the theorem appears in Section 12, and the special values of t_0 in each case are indicated in Table 1.1.

TABLE 1.1. Simply-connected examples with one bubble

Topology of connected sum	Value(s) of t_0
$\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$	$-1/3$
$S^2 \times S^2 \# \overline{\mathbb{CP}}^2 = \mathbb{CP}^2 \# 2\overline{\mathbb{CP}}^2$	$-1/3$, $-(9m_1)^{-1}$
$2\# S^2 \times S^2$	$-2(9m_1)^{-1}$

With $\mathbb{CP}^2$ as a compact factor, there are three fixed points of the torus action, and with $S^2 \times S^2$, there are four fixed points. Employing various discrete symmetries will also allow us to obtain critical metrics on connected sums with more than two factors. Theorem 1.2 extends to the following cases:

- (iv) $3\#S^2 \times S^2$; the product metric on $S^2 \times S^2$ with Green's function $S^2 \times S^2$ metrics attached at two fixed points. In this case, we will impose an additional symmetry called bilateral symmetry.
- (v) $S^2 \times S^2 \# 2\overline{\mathbb{CP}}^2 = \mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$; the product metric on $S^2 \times S^2$ with Burns metrics attached at two fixed points, with bilateral symmetry.
- (vi) $\mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$; the Fubini-Study metric with Burns metrics attached at all fixed points, with a symmetry called trilateral symmetry.
- (vii) $\mathbb{CP}^2 \# 3(S^2 \times S^2) = 4\mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$; the Fubini-Study metric with Green's function $S^2 \times S^2$ metrics attached at all fixed points, with trilateral symmetry.
- (viii) $S^2 \times S^2 \# 4\overline{\mathbb{CP}}^2 = \mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$; the product metric on $S^2 \times S^2$ with Burns metrics attached at all fixed points, with a symmetry called quadrilateral symmetry.
- (ix) $5\#S^2 \times S^2$ viewed as the product metric on $S^2 \times S^2$ with Greens function $S^2 \times S^2$ metrics attached at all fixed points, with quadrilateral symmetry.

The special values of t_0 in each case are indicated in Table 1.2.

Table 1.2. Simply-connected examples with several bubbles

Topology of connected sum	Value of t_0	Symmetry
$3\#S^2 \times S^2$	$-2(9m_1)^{-1}$	bilateral
$S^2 \times S^2 \# 2\overline{\mathbb{CP}}^2 = \mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$	$-1/3$	bilateral
$\mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$	$-1/3$	trilateral
$\mathbb{CP}^2 \# 3(S^2 \times S^2) = 4\mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$	$-(9m_1)^{-1}$	trilateral
$S^2 \times S^2 \# 4\overline{\mathbb{CP}}^2 = \mathbb{CP}^2 \# 5\overline{\mathbb{CP}}^2$	$-1/3$	quadrilateral
$5\#S^2 \times S^2$	$-2(9m_1)^{-1}$	quadrilateral

Remark 1.3. Since $S^2 \times S^2$ admits an orientation-reversing diffeomorphism, there is only one possibility for a connect sum with $S^2 \times S^2$, which is why $\overline{S^2 \times S^2}$ does not appear in the list of examples.

1.4. **Non-simply-connected examples.** The product metric on $S^2 \times S^2$ admits the Einstein quotient $S^2 \times S^2/\mathbb{Z}_2$, where $\mathbb{Z}_2$ acts by the antipodal map on both factors, and the quotient $\mathbb{RP}^2 \times \mathbb{RP}^2$. Using one of these metrics as the compact factor or the Green's function metric of one of these as one of the AF spaces, we can obtain several non-simply-connected examples. We will denote

$$m_2 = \mathrm{mass}(\hat{g}_{S^2 \times S^2/\mathbb{Z}^2}), \tag{1.21}$$

and

$$m_3 = \mathrm{mass}(\hat{g}_{\mathbb{RP}^2 \times \mathbb{RP}^2}). \tag{1.22}$$

Again, by the positive mass theorem, $m_2 > 0$ and $m_3 > 0$. Theorem 1.2 holds for these examples as well, and the special values of t_0 in each non-simply-connected case with one bubble are indicated in Table 1.3. We note those without an $\mathbb{RP}^2 \times \mathbb{RP}^2$

factor are orientable, and those with an $\mathbb{RP}^2 \times \mathbb{RP}^2$ factor are non-orientable. Also note that the first, second, fifth and sixth examples have finite fundamental groups. The others have infinite fundamental group (in particular, by the Myers Theorem these manifolds do not admit positive Einstein metrics).

Table 1.3. Non-simply-connected examples with one bubble

Topology of connected sum	Value(s) of t_0
$(S^2 \times S^2/\mathbb{Z}_2)\#\overline{\mathbb{CP}}^2$	$-1/3$, $-(9m_2)^{-1}$
$(S^2 \times S^2/\mathbb{Z}_2)\#S^2 \times S^2$	$-2(9m_1)^{-1}$, $-2(9m_2)^{-1}$
$(S^2 \times S^2/\mathbb{Z}_2)\#(S^2 \times S^2/\mathbb{Z}_2)$	$-2(9m_2)^{-1}$
$(S^2 \times S^2/\mathbb{Z}_2)\#\mathbb{RP}^2 \times \mathbb{RP}^2$	$-2(9m_3)^{-1}$, $-2(9m_2)^{-1}$
$\mathbb{RP}^2 \times \mathbb{RP}^2\#\overline{\mathbb{CP}}^2$	$-1/3$, $-(9m_3)^{-1}$
$\mathbb{RP}^2 \times \mathbb{RP}^2\#S^2 \times S^2$	$-2(9m_1)^{-1}$, $-2(9m_3)^{-1}$
$\mathbb{RP}^2 \times \mathbb{RP}^2\#\mathbb{RP}^2 \times \mathbb{RP}^2$	$-2(9m_3)^{-1}$

As in the simply-connected case, we can take advantage of various symmetries to obtain non-simply-connected examples with more than one bubble. For the complete list, see Appendix B.

1.5. **The Bach-flat case.** We remark that Theorem 1.1 holds in the Bach-flat case ($t = 0$), provided one restricts to traceless tensors throughout the argument (this is necessary due to conformal invariance of the Bach tensor). This expansion cannot be directly used to produce Bach-flat metrics, since the freedom to move the parameter t is crucial in the proof of Theorem 1.2. However, the main argument does imply the following *non-existence* result:

Theorem 1.4. *Assume that both (Z, g_Z) and (Y, g_Y) are Bach-flat, toric, and admit a diagonal symmetry. Let $z_0 \in Z$ and $y_0 \in Y$ be fixed points of the respective torus actions. If*

$$W(y_0) \circledast W(z_0) \neq 0, \tag{1.23}$$

then there is no equivariant Bach-flat metric in a $C^{4,\alpha}$-neighborhood of the approximate metric.

This is applicable to all of the above examples, so we may conclude that there is no Bach-flat metric near the metrics found in Theorem 1.2. In particular, these metrics are not Einstein. We remark that this non-existence theorem is true without the equivariance assumption, but a complete proof of this adds considerable technical details, so is not included.

Note that in the case of $\mathbb{CP}^2\#\mathbb{CP}^2$, it is easy to see that $W(y_0) \circledast W(z_0) = 0$, since there is an orientation-reversal required when performing the connected sum. This is not surprising, since it is well-known that there is a 1-parameter family of self-dual metrics (which are Bach-flat) near the approximate metric [Poo86, LeB91, Via10].

1.6. **Remarks.** The proof of Theorem 1.2 shows the following dichotomy: either (i) there is a critical metric at exactly the critical t_0, in which case there would necessarily be a 1-dimensional moduli space of solutions for this fixed t_0 (as pointed out above, this indeed happens for $\mathbb{CP}^2\#\mathbb{CP}^2$, in which case there is a 1-parameter family of self-dual metrics). The other possibility (ii) is that for each value of the gluing parameter a sufficiently small, there will be a critical metric for a corresponding value of $t_0 = t_0(a)$. The dependence of t_0 on a will depend on the next term in the expansion of (1.18). For example, if this expansion were improved to

$$\lambda_1 = \lambda a^4 + \mu a^8 + O(a^{12-\epsilon}), \tag{1.24}$$

with $\mu \neq 0$, then we would have the dependence

$$t_0 = \frac{1}{4R(z_0)\mathrm{mass}(g_N)}\Big(- \frac{2}{3}W(y_0) \circledast W(z_0) - \frac{\mu}{\omega_3}a^4\Big) + O(a^{8-\epsilon}). \tag{1.25}$$

as $a \to 0$.

It should be possible to extend the methods in this paper to compute μ. If it turns out that $\mu \neq 0$, then one may conclude that possibility (ii) definitely happens. The sign of μ would then determine if solutions are found for $t > t_0$ or $t < t_0$. If $\mu = 0$, this would indicate (but not prove) that possibility (i) is what actually occurs. The methods in this paper cannot practically be used to determine that possibility (i) actually happens, since there would be an infinite sequence of obstructions to check in this eventuality.

We next make some remarks about some relations between Kähler geometry and the value $t_0 = -1/3$ appearing in the above tables. Using the Hirzebruch signature theorem, we can write

$$\mathcal{B}_{-1/3}[g] = -48\pi^2\sigma(M^4) + 2\int \big(|W^+|^2 - \frac{1}{6}R^2\big)\, dV. \tag{1.26}$$

An immediate corollary of this formula is that if (M^4, g) is Kähler, then

$$\mathcal{B}_{-1/3}[g] = -48\pi^2\sigma(M^4). \tag{1.27}$$

In addition, a constant scalar curvature Kähler metric is necessarily critical for the value $t_0 = -1/3$ [Der83]. We note that important gluing results for constant scalar curvature Kähler metrics were proved in [AP06, AP09].

For the manifolds $\mathbb{CP}^2\#k\overline{\mathbb{CP}}^2$, when $k = 1, 2, 3$, or 5, consider the cases when a Burns metric is used for the bubbles (the cases when $t_0 = -1/3$). In these cases, it is known that there are extremal Kähler metrics near the näive approximate metric [APS11, Szé12]. These extremal metrics do not have constant scalar curvature, so they are not the same as the critical metrics found in Theorem 1.2. There might be some other relation between these metrics (such as conformality), but we are not aware of any such relation. These manifolds are known to admit Einstein metrics [Bes87, CLW08, LeB12].

However, on many of the other manifolds considered in this paper, there does not exist *any* Kähler metric (for example $2\#S^2 \times S^2$), and the critical metrics found in

Theorem 1.2 are the first known "canonical" metrics, to the best of the authors' knowledge.

1.7. **Acknowledgements.** This work was done in part at the Institute for Advanced Study in Princeton and at the Institut Henri Poincaré in Paris. Both authors have been partially supported by the NSF; the first author under NSF grant DMS-1206661 and the second author under NSF grant DMS-1105187. The second author was partially supported by the Simons Foundation as a Simons Fellow of Mathematics. Both authors thank the above institutions for their generous support.

The authors would like to thank Olivier Biquard for crucial assistance in completing this work, and to Simon Brendle, Claude LeBrun, Rafe Mazzeo, Frank Pacard, and Karen Uhlenbeck for many helpful remarks.

2. The building blocks

In this section, we will derive metric expansions for the "building blocks" of our gluing procedure; the Fubini-Study metric and product metric on $S^2 \times S^2$. We will also give metric expansions of the associated scalar-flat asymptotically flat metrics, arising from the Green's function of the conformal Laplacian.

The general gluing problem has many degrees of freedom. We will take advantage of various symmetries to reduce eventually to only one degree of freedom. So in this section, we will also describe the various group actions which will be used for an equivariant gluing.

2.1. **The Fubini-Study metric.** Recall that $\mathbb{CP}^2$ is the set of complex projective lines through the origin in $\mathbb{C}^3$. Equivalently, $\mathbb{CP}^2$ is the set of equivalence classes $\{\mathbb{C}^3 \setminus \{0\}\}/\mathbb{C}^*$, where the action of $\mathbb{C}^*$ is defined by, for $\lambda \in \mathbb{C}^*$,

$$[u_0, u_1, u_2] \mapsto [\lambda u_0, \lambda u_1, \lambda u_2].$$

Let $U_i = \{[u_0, u_1, u_2] | u_i \neq 0\}$, for $i = 0, 1, 2$. The Fubini-Study metric is given in U_0 by [KN96]

$$\begin{aligned} g_{FS} &= \frac{\sqrt{-1}}{2} \partial\bar{\partial}(1 + |u_1|^2 + |u_2|^2) \\ &= \frac{(1+|u|^2)(du_1 d\bar{u}_1 + du_2 d\bar{u}_2) - (\bar{u}_1 du_1 + \bar{u}_2 du_2)(u_1 d\bar{u}_1 + u_2 d\bar{u}_2)}{(1+|u|^2)^2}. \end{aligned} \tag{2.1}$$

This extends to an Einstein metric on $\mathbb{CP}^2$ with $Ric(g) = 6g$, and $PU(3)$, the projective unitary group (the unitary group $U(3)$ modulo its center), acts by isometries.

We will consider two sub-actions of this group action. The first is an action of $U(2)$ fixing the point $[1, 0, 0]$. Globally, this action is given by, for $A \in U(2)$,

$$[u_0, u_1, u_2] \mapsto [u_0, A(u_1, u_2)]. \tag{2.2}$$

The point $[1, 0, 0]$ is the only fixed point of this action. In U_0, this action is given by the standard action of $U(2)$ acting on $\mathbb{C}^2$.

The second action is the torus action of the form

$$[u_0, u_1, u_2] \mapsto [u_0, e^{\sqrt{-1}\theta_1} u_1, e^{\sqrt{-1}\theta_2} u_2] \tag{2.3}$$

where $\theta_1, \theta_2 \in [0, 2\pi]$. This action has 3 fixed points $[1, 0, 0], [0, 1, 0]$, and $[0, 0, 1]$. In U_0, this action is given by

$$(u_1, u_2) \mapsto (e^{\sqrt{-1}\theta_1} u_1, e^{\sqrt{-1}\theta_2} u_2). \tag{2.4}$$

Next, let $\{\sigma_1, \sigma_2, \sigma_3\}$ be a left-invariant coframing on S^3 such that σ_3 is a connection form for the Hopf fibration $\pi : S^3 \to S^2 = \mathbb{CP}^1$ defined by

$$\pi(u_1, u_2) = [u_1 : u_2], \tag{2.5}$$

and such that $\pi^* g_{S^2} = 4(\sigma_1^2 + \sigma_2^2)$. The Fubini-Study metric can then be written as [EGH80, page 257]

$$g_{FS} = \frac{1}{(1+r^2)^2} dr^2 + \frac{r^2}{1+r^2}\Big(\sigma_1^2 + \sigma_2^2 + \frac{1}{1+r^2}\sigma_3^2\Big) \tag{2.6}$$

From this expression, the above action of $U(2)$ is seen here here as an action of $\mathrm{SO}(3) \times \mathrm{SO}(2)$ where the first factor acts by rotations of S^2, and the second factor acts by rotations of the fiber of the Hopf fibration. The above torus action is the restricted action where the first factor acts by a rotation of S^2 fixing the north and south pole.

From (2.6), we see that $\rho = \arctan(r)$ is the geodesic distance from the basepoint, and under this radial change of coordinates the metric is written as [LNN97]

$$g_{FS} = d\rho^2 + \sin^2(\rho)\big(\sigma_1^2 + \sigma_2^2 + \cos^2(\rho)\sigma_3^2\big), \tag{2.7}$$

with the restriction that $0 < \rho < \pi/2$. Since the coordinate change is radial, we note the important fact that in these coordinates, the above action of $U(2)$ is still the standard linear action.

Finally, we let $\{z^i\}$ be Euclidean normal coordinates, based at $[1, 0, 0]$, so that $U(2)$ acts linearly, and that the above torus action acts by

$$(z_1, z_2, z_3, z_4) \mapsto \big(e^{\sqrt{-1}\theta_1}(z_1 + \sqrt{-1}z_2), e^{\sqrt{-1}\theta_2}(z_3 + \sqrt{-1}z_4)\big). \tag{2.8}$$

In this coordinate system, we have the expansion

$$g_{ij} = \delta_{ij} - \frac{1}{3} R_{ikjl}([1, 0, 0]) z^k z^l + O^{(4)}(|z|^4)_{ij} \tag{2.9}$$

as $|z| \to 0$.

Remark 2.1. We adopt the convention that for a function (or tensor) $f = f(z)$, $f = O^{(m)}(|z|^\alpha)$ means $|\partial^k f| = O(|z|^{\alpha-k})$ for all $1 \le k \le m$ (as z approaches an indicated limit).

This metric is invariant under the diagonal symmetry:

$$(z_1, z_2, z_3, z_4) \mapsto (z_3, z_4, z_1, z_2), \tag{2.10}$$

which is contained in $U(2)$. In the case of toric invariance, we will impose this as an extra symmetry for the equivariant gluing problem. In both cases, there will therefore be only one fixed point, the point $[1,0,0]$. These symmetries are illustrated in Figure 2.1.



FIGURE 2.1. Orbit space of the torus action on $\mathbb{CP}^2$. The vertices of the triangle are fixed points, open edge points are circle orbits, and interior points are principal orbits. The diagonal symmetry is a reflection in the dotted diagonal line passing through [1,0,0]. Invariance under reflection in all dotted diagonal lines will be called trilateral symmetry (note these reflections correspond to coordinate flips $u_i \leftrightarrow u_j$ on $\mathbb{CP}^2$, which are isometries of g_{FS}).

2.2. **The Burns metric.** We begin with a general result regarding the Green's function expansion for a toric Einstein manifold:

Proposition 2.2. *Let G be the Green's function for the conformal Laplacian at the point $p \in M$, where (M,g) is an Einstein metric with positive scalar curvature. If (M,g) admits a non-trivial torus action fixing the point p, then in a Riemannian normal coordinate system $\{z^i\}$, we have the following expansion: For any $\epsilon > 0$,*

$$G = |z|^{-2} + A + O^{(4)}(|z|^{2-\epsilon}) \tag{2.11}$$

as $|z| \to 0$, where A is a constant (independent of ϵ).

Proof. A straightforward computation, which we omit, shows that there is a formal power series solution of the form with leading terms

$$G = |z|^{-2} + A + \dots, \tag{2.12}$$

and A is a constant. Recall that the indicial roots of the Laplacian are $\mathbb{Z} \setminus \{-1\}$. Solutions corresponding to the indicial root 1 are linear, and not invariant under the torus action, so there is no linear term in the expansion. It follows from standard techniques that the formal expansion (2.12) implies the actual expansion (2.11). The proof is identical to [LP87, Lemma 6.4] (using Riemannian normal coordinates instead of conformal normal coordinates), so the details are omitted. □

In the case of the Fubini-Study metric, we have the following improved expansion:

Proposition 2.3. *Let G be the Green's function for the conformal Laplacian of the Fubini-Study metric based at $[1,0,0]$, normalized so that $Ric(g) = 6g$. Then in the above normal coordinate system $\{z^i\}$ we have the expansion*

$$G = |z|^{-2} + \frac{1}{3} + O^{(4)}(|z|^2) \tag{2.13}$$

as $|z| \to 0$.

Proof. Since the metric is invariant under $U(2)$, from uniqueness of the Green's function, G must be radial. Using that $R = 24$, the equation is

$$\Delta G = 4G. \tag{2.14}$$

We let $\rho = |z|$ denote the radial distance function. For a radial function, (2.14) reduces to the ODE

$$G_{\rho\rho} + (3\cot(\rho) - \tan(\rho))G_\rho = 4G \tag{2.15}$$

on the interval $[0, \pi/2]$. This ODE has the general solution

$$G = \frac{C_1}{\sin^2(\rho)} + C_2 \frac{\log(\cos(\rho))}{\sin^2(\rho)} \tag{2.16}$$

for constants C_1 and C_2. The boundary condition $G = \rho^{-2}(1 + o(1))$ as $\rho \to 0$ implies that $C_1 = 1$. For the other boundary condition, in order to give a smooth global solution, we require that $G_\rho(\pi/2) = 0$, which implies that $C_2 = 0$. The claimed expansion follows easily from

$$(\sin\rho)^{-2} = \rho^{-2} + \frac{1}{3} + \frac{1}{15}\rho^2 + O^{(4)}(\rho^4) \tag{2.17}$$

as $\rho \to 0$. □

Since g_{FS} is Bach-flat (it is self-dual with respect to the complex orientation), the metric $g_N = G^2 g_{FS}$ is also Bach-flat, and scalar-flat. Consequently, g_N is B^t-flat for any $t \in \mathbb{R}$. Let $\{x^i = z^i/|z|^2\}$ denote inverted normal coordinates near $[1,0,0]$,and let

$$\mathcal{I}(x) = \frac{x}{|x|^2} = z \tag{2.18}$$

denote the inversion map. With respect to these coordinates, we can write the metric g_N in the complement of a large ball as

$$\begin{aligned} g_N &= \mathcal{I}^*(G^2 g_{FS}) \\ &= (G\circ\mathcal{I})^2\mathcal{I}^*\Big(\{\delta_{ij} - \frac{1}{3}R_{ikjl}([1,0,0])z^kz^l + O^{(4)}(|z|^4)_{ij}\}dz^idz^j\Big) \\ &= (|x|^2 + A + O^{(4)}(|x|^{-2}))^2\{\delta_{ij} - \frac{1}{3}R_{ikjl}([1,0,0])\frac{x^kx^l}{|x|^4} + O^{(4)}(|x|^{-4})_{ij}\} \\ &\quad \cdot \frac{1}{|x|^2}\Big(\delta_{ip} - \frac{2}{|x|^2}x^ix^p\Big)dx^p \cdot \frac{1}{|x|^2}\Big(\delta_{jq} - \frac{2}{|x|^2}x^jx^q\Big)dx^q, \end{aligned} \tag{2.19}$$

so we have the expansion

$$(g_N)_{ij}(x) = \delta_{ij} - \frac{1}{3}R_{ikjl}([1,0,0])\frac{x^kx^l}{|x|^4} + 2A\frac{1}{|x|^2}\delta_{ij} + O^{(4)}(|x|^{-4}) \tag{2.20}$$

as $|x| \to \infty$. Clearly, g_N is asymptotically flat (AF) of order $\gamma = 2$.

Note that this metric is also invariant under the standard linear action of $U(2)$, now acting in the $\{x\}$-coordinates.

Remark 2.4. As the title of the subsection indicates, this metric is also known as the Burns metric; it is a Kähler scalar-flat metric on the blow-up of $\mathbb{C}^2$ at the origin. By the coordinate change $r = \sin^{-1}(\rho)$, and multiplying by r^4, one obtains

$$g_N = \frac{dr^2}{1 - r^{-2}} + r^2\Big[\sigma_1^2 + \sigma_2^2 + (1 - r^{-2})\sigma_3^2\Big], \tag{2.21}$$

which is the expression of the Burns metric obtained in [LeB88]. We could instead use this coordinate system for the Burns metric in this paper. However, since there is not an analogue of this for the next example, we will remain with the above inverted Riemannian normal coordinates, in order to give a unified approach.

We note here the following, which relates the constant A to the mass of the Green's function metric, and will be used later.

Proposition 2.5. *Let (M, g) be as in Proposition 2.2. Then the mass of the AF metric $g_N = G^2g$ on $N = M \setminus \{p\}$ is given by*

$$\text{mass}(g_N) = 12A - \frac{R(p)}{12}. \tag{2.22}$$

Proof. This follows from (1.15) using inverted normal coordinates; the routine calculation is omitted. ☐

For the Fubini-Study metric, since $R(p) = 24$, and $A = 1/3$, this implies that

$$\text{mass}(g_N) = 2. \tag{2.23}$$

2.3. **The product metric on $S^2 \times S^2$.** Next, we consider $S^2 \times S^2$ with metric $g = g_{S^2} \times g_{S^2}$ the product of metrics of constant Gaussian curvature 1. The torus action we will consider is just the product of counter-clockwise S^1-rotations fixing the north and south poles. This action has 4 fixed points $(n,n),(n,s),(s,n)$, and (s,s), where n and s are the north and south poles, respectively.

Taking normal coordinates on each factor around $(n,n) \in S^2 \times S^2$, yields a normal coordinate system $(r_1,\theta_1,r_2,\theta_2)$ so that

$$g_{S^2\times S^2} = dr_1^2 + \sin^2(r_1)d\theta_1^2 + dr_2^2 + \sin^2(r_2)d\theta_2^2, \tag{2.24}$$

and the radial distance function is given by $\rho = \sqrt{r_1^2+r_2^2}$. Finally, we let $\{z^i\}$ be Euclidean normal coordinates based at (n,n), so that the above torus action acts by

$$(z_1,z_2,z_3,z_4) \mapsto \big(e^{\sqrt{-1}\theta_1}(z_1+\sqrt{-1}z_2), e^{\sqrt{-1}\theta_2}(z_3+\sqrt{-1}z_4)\big). \tag{2.25}$$

In this coordinate system, we have the expansion

$$g_{ij} = \delta_{ij} - \frac{1}{3}R_{ikjl}((n,n))z^kz^l + O(|z|^4)_{ij}, \tag{2.26}$$

as $|z| \to 0$.

In addition to toric invariance, this metric is also invariant under the diagonal symmetry:

$$(z_1,z_2,z_3,z_4) \mapsto (z_3,z_4,z_1,z_2). \tag{2.27}$$

We will also impose this as an extra symmetry for the equivariant gluing problem. These symmetries, as well as some other symmetries we will use later, are illustrated in Figure 2.2.

As mentioned in the introduction, the product metric on $S^2\times S^2$ admits the Einstein quotient $S^2 \times S^2/\mathbb{Z}_2$, where $\mathbb{Z}_2$ acts by the antipodal map on both factors, and the quotient $\mathbb{RP}^2 \times \mathbb{RP}^2$. These quotients are also toric and the same expansion (2.26) holds for these. The diagonal symmetry also descends to a symmetry of these metrics.

2.4. **Green's function of product metric.** Let G be the Green's function for the conformal Laplacian of the product metric at the point (n,n), normalized so that $R = 4$. By Proposition 2.2, in the above normal coordinate system $\{z^i\}$, for any $\epsilon > 0$, we have the expansion

$$G = |z|^{-2} + A + O^{(4)}(|z|^{2-\epsilon}) \tag{2.28}$$

as $|z| \to 0$, where A is a constant (independent of ϵ).

Since $g_{S^2\times S^2}$ is Bach-flat (it is Einstein), the metric $g_N = G^2 g_{S^2\times S^2}$ is also Bach-flat, and scalar-flat. Consequently, g_N is B^t-flat for any $t \in \mathbb{R}$. Letting $\{x^i = z^i/|z|^2\}$ denote inverted normal coordinates, analogous to (2.20), the metric g_N admits the expansion

$$(g_N)_{ij}(x) = \delta_{ij} - \frac{1}{3}R_{ikjl}((n,n))\frac{x^kx^l}{|x|^4} + 2A\frac{1}{|x|^2}\delta_{ij} + O^{(4)}(|x|^{-4+\epsilon}) \tag{2.29}$$

as $|x| \to \infty$, for any $\epsilon > 0$. Clearly, g_N is AF of order $\gamma = 2$.



FIGURE 2.2. Orbit space of the torus action on $S^2 \times S^2$. The vertices of the square are fixed points, open edge points are circle orbits, and interior points are principal orbits. The diagonal symmetry is a reflection in the dotted diagonal line passing through (n, n). The bilateral symmetry is reflection in the dotted anti-diagonal line. Reflection in the dotted vertical line is the antipodal map of the first factor, while reflection in the dotted horizontal line is the antipodal map of the second factor. Invariance under all of these reflections will be called quadrilateral symmetry.

This metric is invariant under the above diagonal torus action, now acting in the $\{x\}$-coordinates, and is also invariant under the diagonal symmetry

$$(x_1, x_2, x_3, x_4) \mapsto (x_3, x_4, x_1, x_2). \tag{2.30}$$

Remark 2.6. Unlike the case of the Burns metric, there is no explicit description of this metric known (to the best of the authors' knowledge). Since the metric is invariant under the above torus action, from uniqueness of the Green's function, $G = G(r_1, r_2)$. Using that $R = 4$, the equation is

$$\Delta G = \frac{2}{3} G. \tag{2.31}$$

Since $G = G(r_1, r_2)$, a computation shows that this reduces to the PDE

$$G_{r_1 r_1} + \cot(r_1) G_{r_1} + G_{r_2 r_2} + \cot(r_2) G_{r_2} = \frac{2}{3} G, \tag{2.32}$$

on the square $[0, \pi] \times [0, \pi]$. Unlike the case of the Fubini-Study metric, this does not appear to admit any explicit solution.

3. The nonlinear map

Let (M, g) be a compact manifold of dimension 4, and Let $S^2(T^*M)$ denote the bundle of symmetric 2-tensors on M. We recall some important linear operators. For simplicity of notation, we will treat the domain and range of an operator as if it were the bundle itself, although the operator really acts on sections of the bundle. Let $\delta_g : S^2(T^*M) \to T^*M$ denote the divergence operator

$$(\delta_g h)_j = \nabla^i h_{ij}, \tag{3.1}$$

and $\delta^* : T^*M \to S^2(T^*M)$ its L^2-adjoint. Note that

$$\delta^* = -\frac{1}{2}\mathcal{L}, \tag{3.2}$$

where $\mathcal{L}$ is the Killing operator:

$$(\mathcal{L}_g \omega)_{ij} = \nabla_i \omega_j + \nabla_j \omega_i. \tag{3.3}$$

We let $\mathcal{K}_g$ denote the conformal Killing operator, the trace-free part of $\mathcal{L}_g$:

$$(\mathcal{K}_g \omega)_{ij} = \nabla_i \omega_j + \nabla_j \omega_i - \frac{1}{2}(\delta_g \omega) g_{ij}. \tag{3.4}$$

Next, for a fixed background metric g, define the nonlinear map P_g

$$P_g^t : C^{4,\alpha}(S^2(T^*M)) \to C^{0,\alpha}(S^2(T^*M)) \tag{3.5}$$

by

$$P_g^t(\theta) = B^t(g + \theta) + \mathcal{K}_{g+\theta}\delta_g \mathcal{K}_g \delta_g \overset{\circ}{\theta}, \tag{3.6}$$

where

$$\overset{\circ}{\theta} = \theta - \frac{1}{4} tr_g(\theta) g. \tag{3.7}$$

Remark 3.1. The domain of P_g^t is not actually the entire space; it is the subset of $C^{4,\alpha}$ so that $g + \theta$ is a Riemannian metric. The fact that the image lies in $C^{0,\alpha}$ is a consequence of P_g^t being analytic as a function of θ and its derivatives up to order four.

We let $S_g^t \equiv (P_g^t)'(0)$ denote the linearized operator at $\theta = 0$.

Remark 3.2. When the base metric is clear from the context, we will often omit the subscript in the operator P^t and its linearization S^t. To further simplify notation, we will also often omit the superscript t from both of these operators, since it is clear that they depend on t.

Proposition 3.3. *If $t \neq 0$, then S^t is elliptic.*

Proof. This is proved in [GV11, Theorem 2.7 (i)], although we provide a brief sketch since some of the formulas will be needed in subsequent sections. We also note a difference in notation with our previous paper [GV11]. In that paper we considered the functional

$$\mathcal{F}_\tau = \int_M |Ric|^2 dV + \tau \int_M R^2 dV. \tag{3.8}$$

From (1.5), we obtain the relation

$$\mathcal{F}_\tau = 16\pi^2\chi(M) + \frac{1}{2}\mathcal{B}_{2(\tau+\frac{1}{3})}. \tag{3.9}$$

Taking gradients, we obtain the relation

$$\nabla \mathcal{B}_t = 2\nabla \mathcal{F}_{\frac{t}{2}-\frac{1}{3}}. \tag{3.10}$$

It follows from the formula for P that the linearized operator is given by

$$S^t h = (B' + tC')h + \mathcal{K}_g\delta_g\mathcal{K}_g\delta_g\overset{\circ}{h}, \tag{3.11}$$

where B' and C' are the linearizations of B and C respectively. Using (3.10), from [GV11, Equation (2.54)] the leading terms of $B' + tC'$ are

(3.12)

$$\begin{aligned}(B'+tC')h_{ij} = \Delta^2 h_{ij} - \Delta\big[\nabla_i\delta_j h + \nabla_j\delta_i h\big] - (2t - \frac{1}{3})\nabla_i\nabla_j(\Delta tr\ h)\\ + (2t + \frac{2}{3})\nabla_i\nabla_j(\delta^2 h) + (2t - \frac{1}{3})\big[\Delta^2(tr\ h) - \Delta(\delta^2 h)\big]g_{ij} + \cdots .\end{aligned}$$

Also, a simple calculation gives

$$\begin{aligned}(\mathcal{K}_g\delta_g\mathcal{K}_g\delta_g\overset{\circ}{h})_{ij} = \Delta\big[\nabla_i\delta_j h + \nabla_j\delta_i h\big] - \frac{3}{4}\nabla_i\nabla_j(\Delta tr\ h) + \nabla_i\nabla_j(\delta^2 h)\\ + \frac{3}{16}\Delta^2(tr\ h)g_{ij} - \frac{3}{4}\Delta(\delta^2 h)g_{ij} + \cdots .\end{aligned}$$

Consequently,

$$\begin{aligned}Sh = \Delta^2 h - 2\big(t + \frac{5}{24}\big)\nabla^2(\Delta tr\ h) - 2\big(t + \frac{5}{24}\big)\Delta(\delta^2 h)g\\ + 2\big(t + \frac{5}{6}\big)\nabla^2(\delta^2 h) + 2\big(t - \frac{7}{96}\big)\Delta^2(tr\ h)g + \cdots .\end{aligned} \tag{3.13}$$

It follows from (3.13) that the symbol of S is

$$\begin{aligned}(\sigma_\xi S)h_{ij} = |\xi|^4 h - 2\big(t + \frac{5}{24}\big)\xi_i\xi_j|\xi|^2(tr\ h) - 2\big(t + \frac{5}{24}\big)|\xi|^2 h_{k\ell}\xi_k\xi_\ell g_{ij}\\ + 2\big(t + \frac{5}{6}\big)\xi_i\xi_j h_{k\ell}\xi_k\xi_\ell + 2\big(t - \frac{7}{96}\big)|\xi|^4(tr\ h)g_{ij},\end{aligned} \tag{3.14}$$

which is elliptic for $t \neq 0$, according to [GV11, Theorem 2.7 (i)]. □

Remark 3.4. For purposes below, it will be useful to rewrite (3.13) as

$$Sh = \Delta^2[h - \frac{1}{4}(tr\ h)g] + \mathcal{K}\big[d(\mathcal{D}_2(h))\big] + \frac{3}{2}t\Big[\Delta^2(tr\ h) - \Delta(\delta^2 h)\Big]g + \cdots \tag{3.15}$$

where $\mathcal{D}_2 : S^2T^*M \to C^\infty$ is a second-order operator given by

$$\mathcal{D}_2(h) = \big(t + \frac{5}{6}\big)\delta^2 h - \big(t + \frac{5}{24}\big)\Delta(tr\ h). \tag{3.16}$$

The following proposition shows that the zeroes of P are in fact B^t-flat metrics:

Proposition 3.5. *Assume* $t \neq 0$. *If* $P(\theta) = 0$ *and* $\theta \in C^{4,\alpha}$ *for some* $0 < \alpha < 1$, *then* $B^t(g+\theta) = 0$ *and* $\theta \in C^\infty$.

Proof. The equation is

$$B^t(g+\theta) + \mathcal{K}_{g+\theta}\delta_g\mathcal{K}_g\delta_g\overset{\circ}{\theta} = 0. \tag{3.17}$$

We claim that both terms on the left hand side of (3.17) vanish. The proof involves an integration by parts argument, but this presents a difficulty since $\theta \in C^{4,\alpha}$ only implies that $P_g(\theta)$ is C^α, and not necessarily differentiable. To get around this problem we mollify θ; i.e., let $\{\theta_\epsilon\}$ be a family of smooth tensor fields such that $\theta_\epsilon \to \theta$ in $C^{4,\alpha}$ as $\epsilon \to 0$, and let $\tilde{g}_\epsilon = g + \theta_\epsilon$. From (3.17) and the continuity of P it follows that

$$\eta_\epsilon = B^t(g+\theta_\epsilon) + \mathcal{K}_{g+\theta_\epsilon}[\Box_{\mathcal{K}_g}\beta_g\theta_\epsilon], \tag{3.18}$$

where $\eta_\epsilon \to 0$ in C^α. Pair both sides of (3.18) with $\mathcal{L}_{\tilde{g}_\epsilon}[\Box_{\mathcal{K}_g}\beta_g\theta_\epsilon]$ (with respect to the L^2-inner product defined by $\tilde{g}_\epsilon$), where $\mathcal{L}$ is the Killing operator defined in (3.3):

$$\begin{aligned}\langle \mathcal{L}_{\tilde{g}_\epsilon}[\Box_{\mathcal{K}_g}\beta_g\theta_\epsilon], \eta_\epsilon\rangle_{L^2} &= \big\langle \mathcal{L}_{\tilde{g}_\epsilon}[\Box_{\mathcal{K}_g}\beta_g\theta_\epsilon], B^t(\tilde{g}_\epsilon) + \mathcal{K}_{\tilde{g}_\epsilon}[\Box_{\mathcal{K}_g}\beta_g\theta_\epsilon]\big\rangle_{L^2} \\ &= \big\langle \mathcal{L}_{\tilde{g}_\epsilon}[\Box_{\mathcal{K}_g}\beta_g\theta_\epsilon], B^t(\tilde{g}_\epsilon)\big\rangle_{L^2} + \|\mathcal{K}_{\tilde{g}_\epsilon}[\Box_{\mathcal{K}_g}\beta_g\theta_\epsilon]\|^2_{L^2}.\end{aligned}$$

Integrating by parts in the first term on the right-hand side, we get

$$\big\langle \mathcal{L}_{\tilde{g}_\epsilon}[\Box_{\mathcal{K}_g}\beta_g\theta_\epsilon], B^t(\tilde{g}_\epsilon)\big\rangle_{L^2} = -2\big\langle \Box_{\mathcal{K}_g}\beta_g\theta_\epsilon, \delta_{\tilde{g}_\epsilon}\big(B^t(\tilde{g}_\epsilon)\big)\big\rangle_{L^2} = 0,$$

since $\mathcal{L}^* = -2\delta$ and the gradient of a Riemannian functional is always divergence-free (see [Bes87], Proposition 4.11). Therefore,

$$\langle \mathcal{L}_{\tilde{g}_\epsilon}[\Box_{\mathcal{K}_g}\beta_g\theta_\epsilon], \eta_\epsilon\rangle_{L^2} = \frac{1}{2}\|\mathcal{K}_{\tilde{g}_\epsilon}[\Box_{\mathcal{K}_g}\beta_g\theta_\epsilon]\|^2_{L^2}.$$

Letting $\epsilon \to 0$, the left-hand side converges to zero, while the right-hand side converges to $\mathcal{K}_{\tilde{g}}[\Box_{\mathcal{K}_g}\beta_g\theta]$, which consequently vanishes. We conclude that

$$B^t(\tilde{g}) = 0 \tag{3.19}$$

as claimed.

Next, taking a trace of (3.19), yields

$$-6t\Delta R_{\tilde{g}} = 0, \tag{3.20}$$

which implies that the scalar curvature of $\tilde{g}$ is constant. The equation (3.19) then implies that $\Delta_{\tilde{g}}Ric_{\tilde{g}} \in C^{2,\alpha}$ (more precisely, around any point $p \in M$, there exists a coordinate system $\{x^i\}$ such that the components are in $C^{2,\alpha}$), which implies that $Ric_{\tilde{g}} \in C^{4,\alpha}$. Since $\tilde{g} \in C^{4,\alpha}$, there exists a harmonic coordinate system $\{y^i\}$ around p such that the equation

$$\frac{1}{2}\tilde{g}^{ij}\partial^2_{ij}\tilde{g}_{kl} + Q_{kl}(\partial\tilde{g}, \tilde{g}) = -Ric_{kl}(\tilde{g}) \tag{3.21}$$

holds, where $Q(\partial\tilde{g},\tilde{g})$ is an expression that is quadratic in $\partial\tilde{g}$, polynomial in $\tilde{g}$ and has $\sqrt{|\tilde{g}|}$ in its denominator [Pet06]. From this we conclude that $\tilde{g}_{ij} \in C^{5,\alpha}$. A bootstrap argument shows that $\tilde{g}_{ij} \in C^{\ell,\alpha}$ for any $\ell > 0$.

$\square$

Later, we will view the nonlinear map in (3.6) as a mapping from

$$P_g : C^{4,\alpha}_{\delta} \to C^{0,\alpha}_{\delta-4}, \tag{3.22}$$

where the spaces are certain weighted Hölder spaces with weight function $w > 0$. Of course, since $w > 0$ and M is compact, these norms are equivalent to the usual Hölder norms. However, in the gluing construction, the weight function will become large, and these norms will then not be uniformly equivalent to the usual norms.

Next, we define the weighted norms we will use. For $\delta \in \mathbb{R}$, and a positive weight function $w > 0$,

$$\|h\|_{C^0_\delta} \equiv \|w^{-\delta}h\|_{C^0} = \sup_{x\in M} |w^{-\delta}(x)h(x)|. \tag{3.23}$$

For $0 < \alpha < 1$, define the semi-norm

$$|h|_{C^{0,\alpha}_\delta} \equiv \sup_{x\in M} \Big(w^{-\delta+\alpha}(x) \sup_{0<4d(x,y)\leq w(x)} \frac{|h(x)-h(y)|}{d(x,y)^\alpha} \Big). \tag{3.24}$$

Finally, define the norm

$$\|h\|_{C^{k,\alpha}_\delta} \equiv \sum_{i=0}^{k} \|\nabla^i h\|_{C^0_{\delta-i}} + |\nabla^k h|_{C^{0,\alpha}_\delta}. \tag{3.25}$$

Remark 3.6. For the remainder of the paper, we fix $\alpha \in \mathbb{R}$ satisyfing $0 < \alpha < 1$.

3.1. **Estimate on the nonlinear terms.**

The following proposition regarding the nonlinear structure of the operator P_g is crucial and will be used througout the paper.

Proposition 3.7. *Write*

$$P_g(h) = P_g(0) + S_g h + Q_g(h), \tag{3.26}$$

where S_g is the linearization of P. Then we have the following:

(i) If $h \in C^{4,\alpha}$ with $\|h\|_{C^0} < s_0$ small, then there exists a constant $C_1 = C_1(s_0)$ so that Q_g satisfies

$$\begin{aligned}|Q_g(h)| \leq C_1\Big\{ &(|\nabla^2 Rm_g| + |Rm_g|^2)|h|^2 + |\nabla Rm_g||h||\nabla^2 h| + |\nabla Rm_g||h||\nabla h| \\ &+ |Rm_g||h||\nabla^2 h| + |Rm_g||h||\nabla h|^2 + |h||\nabla^4 h| \\ &+ |\nabla h||\nabla^3 h| + |\nabla^2 h|^2 + |\nabla h|^2|\nabla^2 h| + |\nabla h|^4 \Big\}.\end{aligned} \tag{3.27}$$

(ii) Let w denote a weight function, and assume

$$\begin{aligned} w &\geq 1, \\ \delta &< 0. \end{aligned} \tag{3.28}$$

In addition, assume there is a constant $C_0 > 0$ such that

$$\begin{aligned} w^2 |Rm_g| &\leq C_0, \\ w^3 |\nabla_g Rm_g| &\leq C_0, \\ w^4 |\nabla_g^2 Rm_g| &\leq C_0. \end{aligned} \tag{3.29}$$

Then, for $h_i \in C^{4,\alpha}_{\delta}$ with $\|h_i\|_{C^{4,\alpha}_{\delta}} < s_0$ small, there exists a constant $C_2 = C_2(s_0)$ so that Q_g satisfies the following estimate:

$$\|Q_g(h_1) - Q_g(h_2)\|_{C^{0,\alpha}_{\delta-4}} \leq C_2(\|h_1\|_{C^{4,\alpha}_{\delta}} + \|h_2\|_{C^{4,\alpha}_{\delta}}) \cdot \|h_1 - h_2\|_{C^{4,\alpha}_{\delta}}. \tag{3.30}$$

Proof. Since the proof involves a rather lengthy calculation we begin with a brief overview. The tensor $B + tC$ can be schematically expressed as

$$B_g + tC_g = g * g^{-1} * g^{-1} * \nabla_g^2 Rm_g + g * g^{-1} * g^{-1} * Rm_g * Rm_g, \tag{3.31}$$

where Rm_g denotes the curvature tensor of g, $g^{-1} * \cdots * g^{-1} * A * B$ denotes any linear combination of terms involving contractions of the tensor product $A \otimes B$, and $g^{-1} * \cdots * g^{-1} * \nabla_g^k * A$ denotes linear combinations of contractions of the k-th iterated covariant derivative of A. Since the mapping P is defined by

$$P_g(h) = B_{g+h} + tC_{g+h} + \mathcal{K}_{g+h} \Box_g \beta_g h, \tag{3.32}$$

the first step in proving the estimates is to analyze the curvature term

$$\begin{aligned} B_{g+h} + tC_{g+h} = (g+h) * (g+h)^{-1} * (g+h)^{-1} * \nabla_{g+h}^2 Rm_{g+h} \\ + (g+h) * (g+h)^{-1} * (g+h)^{-1} * Rm_{g+h} * Rm_{g+h}. \end{aligned} \tag{3.33}$$

The starting point is the formula

$$\Gamma(g+h)_{ij}^k = \Gamma(g)_{ij}^k + \frac{1}{2}(g+h)^{km} \left\{ \nabla_j h_{im} + \nabla_i h_{jm} - \nabla_m h_{ij} \right\}, \tag{3.34}$$

where $\Gamma(\cdot)$ denotes the Christoffel symbols of a metric. In the following, any covariant derivative without a subscript will mean with respect to the fixed metric g. Using this formula and the notation introduced above, we can express the covariant derivative with respect to the metric $g + h$ as

$$\nabla_{g+h} T = \nabla_g T + (g+h)^{-1} * \nabla_g h * T, \tag{3.35}$$

where T is any tensor field. Also, by the standard formula for the $(1,3)$-curvature tensor in terms of the Christoffel symbols we have

$$Rm_{g+h} = Rm_g + (g+h)^{-1} * \nabla^2 h + (g+h)^{-2} * \nabla h * \nabla h. \tag{3.36}$$

Taking the covariant derivative ∇_{g+h} of Rm_{g+h} and repeatedly using (3.35), we obtain

$$\begin{aligned} \nabla_{g+h} Rm_{g+h} = \nabla Rm_g + (g+h)^{-1} * Rm_g * \nabla h + (g+h)^{-1} * \nabla^3 h \\ + (g+h)^{-2} * (\nabla^2 h * \nabla h) + (g+h)^{-3} * (\nabla h * \nabla h * \nabla h). \end{aligned} \tag{3.37}$$

Differentiating again, repeating the above procedure and collecting terms we have

$$\begin{aligned}\nabla^2_{g+h} Rm_{g+h} &= \nabla^2 Rm_g + (g+h)^{-1} * \nabla Rm_g * \nabla h + (g+h)^{-1} * Rm_g * \nabla^2 h\\ &\quad + (g+h)^{-2} * Rm_g * \nabla h * \nabla h + (g+h)^{-1} * \nabla^4_g h\\ &\quad + (g+h)^{-2} * \nabla^3 h * \nabla h + (g+h)^{-2} * \nabla^2 h * \nabla^2 h\\ &\quad + (g+h)^{-3} * \nabla^2 h * \nabla h * \nabla h + (g+h)^{-4} * \nabla h * \nabla h * \nabla h * \nabla h.\end{aligned} \tag{3.38}$$

Therefore,

$$\begin{aligned}&(g+h) * (g+h)^{-2} * \nabla^2_{g+h} Rm_{g+h} =\\ &\quad (g+h) * \Big\{ (g+h)^{-2} * \nabla^2 Rm_g + (g+h)^{-3} * \nabla Rm_g * \nabla h\\ &\quad + (g+h)^{-3} * Rm_g * \nabla^2 h + (g+h)^{-4} * Rm_g * \nabla h * \nabla h\\ &\quad + (g+h)^{-3} * \nabla^4 h + (g+h)^{-4} * \nabla^3 h * \nabla h\\ &\quad + (g+h)^{-4} * \nabla^2 h * \nabla^2 h + (g+h)^{-5} * \nabla^2 h * \nabla h * \nabla h +\\ &\quad + (g+h)^{-6} * \nabla h * \nabla h * \nabla h * \nabla h \Big\}.\end{aligned} \tag{3.39}$$

Using (3.36), we have a similar expression for the second term in (3.33):

$$\begin{aligned}&(g+h) * (g+h)^{-2} * Rm_{g+h} * Rm_{g+h}\\ &= (g+h) * \Big\{ (g+h)^{-2} * Rm_g * Rm_g + (g+h)^{-3} * Rm_g * \nabla^2 h\\ &+ (g+h)^{-4} * Rm_g * \nabla h * \nabla h + (g+h)^{-4} * \nabla^2 h * \nabla^2 h\\ &+ (g+h)^{-5} * \nabla^2 h * \nabla h * \nabla h + (g+h)^{-6} * \nabla h * \nabla h * \nabla h * \nabla h \Big\}.\end{aligned} \tag{3.40}$$

Combining (3.39) and (3.40) gives

$$\begin{aligned}&(g+h) * \big\{ (g+h)^{-2} * \nabla^2_{g+h} Rm_{g+h} + (g+h)^{-2} * Rm_{g+h} * Rm_{g+h} \big\}\\ &= (g+h) * \big\{ (g+h)^{-2} * \nabla^2 Rm_g + (g+h)^{-2} * Rm_g * Rm_g\\ &+ (g+h)^{-3} * \nabla Rm_g * \nabla h + (g+h)^{-3} * Rm_g * \nabla^2 h\\ &+ (g+h)^{-4} * Rm_g * \nabla h * \nabla h + (g+h)^{-3} * \nabla^4 h\\ &+ (g+h)^{-4} * \nabla^3 h * \nabla h + (g+h)^{-4} * \nabla^2 h * \nabla^2 h\\ &+ (g+h)^{-5} * \nabla^2 h * \nabla h * \nabla h + (g+h)^{-6} * \nabla h * \nabla h * \nabla h * \nabla h \Big\}.\end{aligned} \tag{3.41}$$

Returning to the formula (3.32), the gauge-fixing term can be written

$$\begin{aligned}\mathcal{K}_{g+h} \Box_g \beta_g h &= (g+h)^{-1} * (g+h) * \nabla_{g+h} (\Box_g \beta_g h)\\ &= (g+h)^{-1} * (g+h) * (\nabla_g + (g+h)^{-1} * \nabla h) * (\Box_g \beta_g h)\\ &= (g+h)^{-1} * (g+h) * g^{-3} * g * \nabla^4 h\\ &\quad + (g+h)^{-2} * (g+h) * g^{-3} * g * \nabla h * \nabla^3 h.\end{aligned} \tag{3.42}$$

Combining (3.41) and (3.42) we finally have

(3.43)
$$\begin{aligned} P_g(h) = (g{+}h) * \Big\{ &(g+h)^{-2} * \nabla^2 Rm_g + (g+h)^{-2} * Rm_g * Rm_g \\ &+ (g+h)^{-3} * \nabla Rm_g * \nabla h + (g+h)^{-3} * Rm_g * \nabla^2 h \\ &+ (g+h)^{-4} * Rm_g * \nabla h * \nabla h + (g+h)^{-3} * \nabla^4 h \\ &+ (g+h)^{-1} * g^{-3} * g * \nabla^4 h + (g+h)^{-4} * \nabla^3 h * \nabla h \\ &+ (g+h)^{-2} * g^{-3} * g * \nabla^3 h * \nabla h + (g+h)^{-4} * \nabla^2 h * \nabla^2 h \\ &+ (g+h)^{-5} * \nabla^2 h * \nabla h * \nabla h + (g+h)^{-6} * \nabla h * \nabla h * \nabla h * \nabla h \Big\}. \end{aligned}$$

Since we are trying to estimate the remainder terms in the Taylor expansion of $P(h)$, we want to write the above expression in terms of its linearization; i.e.,

$$\begin{aligned} P_g(h) &= P_g(0) + Sh + \cdots \\ &= B_g + tC_g + Sh + \cdots \end{aligned}$$

To do this, we use the identity (which holds for h small)

$$(g+h)^{-1} - g^{-1} = g^{-2} * h + \sum_{k \geq 2} g^{-k-1} * h^k, \tag{3.44}$$

which follows from the usual geometric series formula. Therefore,

$$(g+h_1)^{-1} - (g+h_2)^{-1} = g^{-2} * (h_1 - h_2) + \sum_{k \geq 2} g^{-k-1} * \big(h_1^k - h_2^k\big). \tag{3.45}$$

Each term in the sum in (3.45) can be written

$$g^{-k-1} * h_1^k - g^{-k-1} * h_2^k = g^{-k-1} * (h_1 - h_2) * \sum_{i+j=k-1} h_1^i * h_2^j. \tag{3.46}$$

Therefore, for h small we can write

$$(g+h)^{-1} - g^{-1} = g^{-2} * h + r_1(h), \tag{3.47}$$

where r_1 satisfies

$$|r_1(h_1) - r_1(h_2)| \leq C(g)\big(|h_1| + |h_2|\big)|h_1 - h_2| \tag{3.48}$$

for h_1, h_2 small. In general we can write

$$(g+h)^{-k} - g^{-k} = g^{-k-1} * h + r_k(h), \tag{3.49}$$

where the remainder satisfies

$$|r_k(h_1) - r_k(h_2)| \leq C_k(g)\big(|h_1| + |h_2|\big)|h_1 - h_2|, \tag{3.50}$$

with a similar estimate for the Hölder norm.

We note that, using the restrictions on the weight function assumed in (3.28), the assumption that $\|h_i\|_{C^{4,\alpha}_\delta}$ is small implies that the C^0-norm of h_i is also small, so we are free to employ (3.50) in the following.

Next, we substitute (3.49) into each term of (3.43) involving a power of $(g+h)^{-1}$, then collect all terms which are zeroth order in h (which combine to give $P_g(0)$), those which are linear in h (which combine to give Sh), and those which are higher order in h. For example, consider the term

$$\begin{aligned}(g+h)*(g+h)^{-3}*\nabla^4 h &= (g+h)*(g^{-3}+g^{-4}*h+r_3(h))*\nabla^4 h\\ &= g*g^{-3}*\nabla^4 h + g*g^{-4}*h*\nabla^4 h + g*r_3(h)*\nabla^4 h\\ &\quad + g^{-3}*h*\nabla^4 h + g^{-4}*h*h*\nabla^4 h + r_3(h)*h*\nabla^4 h.\end{aligned}$$

Next, apply (3.49) to each term in (3.43) in a similar fashion, and write the resulting expression as

$$P_g(h) = P_g(0) + Sh + Q(h), \tag{3.51}$$

where Q is

$$\begin{aligned}Q(h) = (g+h)*\Big\{ & r_2(h)*\nabla^2 Rm_g + r_2(h)*Rm_g*Rm_g + g^{-4}*Rm_g*h*\nabla^2 h\\ & + g^{-4}*\nabla Rm_g*h*\nabla h + r_3(h)*\nabla Rm_g*\nabla h + g^{-4}*Rm_g*h*\nabla^2 h\\ & + r_3(h)*Rm_g*\nabla^2 h + g^{-5}*Rm_g*h*\nabla h*\nabla h\\ & + r_4(h)*Rm_g*\nabla h*\nabla h + g^{-4}*h*\nabla^4 h + g^{-5}*g*h*\nabla^4 h\\ & + g^{-3}*g*r_1(h)*\nabla^4 h + r_3(h)*\nabla^4 h\\ & + g^{-4}*\nabla^3 h*\nabla h + g^{-5}*h*\nabla h*\nabla^3 h + r_4(h)*\nabla h*\nabla^3 h\\ & + g^{-5}*g*\nabla^3 h*\nabla h + g^{-6}*g*h*\nabla h*\nabla^3 h + g^{-3}*g*r_2(h)*\nabla h*\nabla^3 h\\ & + g^{-4}*\nabla^2 h*\nabla^2 h + g^{-5}*h*\nabla^2 h*\nabla^2 h + r_4(h)*\nabla^2 h*\nabla^2 h\\ & + g^{-3}*\nabla h*\nabla h*\nabla^2 h + g^{-6}*h*\nabla h*\nabla h*\nabla^2 h\\ & + r_5(h)*\nabla h*\nabla h*\nabla^2 h + g^{-6}*\nabla h*\nabla h*\nabla h*\nabla h\\ & + g^{-7}*h*\nabla h*\nabla h*\nabla h*\nabla h + r_6(h)*\nabla h*\nabla h*\nabla h*\nabla h\Big\}.\end{aligned} \tag{3.52}$$

The estimate (3.27) follows from considering each term in (3.52), inequality (3.50), and the smallness of h.

We can then prove (3.30) by a fairly straightforward—but, due to the number of terms, very lengthy—process. We will provide the details for estimating some representative terms; the rest can be handled similarly.

For example, consider the term

$$T(h) = g*g^{-4}*h*\nabla^4 h. \tag{3.53}$$

Then

$$\begin{aligned}T(h_1)-T(h_2) &= g*g^{-4}*h_1*\nabla^4 h_1 - g*g^{-4}*h_2*\nabla^4 h_2\\ &= g*g^{-4}*(h_1-h_2)*\nabla^4 h_1 + g*g^{-4}*h_2*\nabla^4(h_1-h_2)\end{aligned}$$

If w denotes the weight, then this implies

$$\begin{aligned}|T(h_1)-T(h_2)|w^{4-\delta} &\leq |h_1-h_2||\nabla^4 h_1|w^{4-\delta}+|h_2||\nabla^4(h_1-h_2)|w^{4-\delta}\\ &= \big\{|h_1-h_2|w^{-\delta}\big\}\big\{|\nabla^4 h_1|w^{4-\delta}\big\}w^{\delta}\\ &\quad+\big\{|h_2|w^{-\delta}\big\}\big\{|\nabla^4(h_1-h_2)|w^{4-\delta}\big\}w^{\delta}.\end{aligned}$$

Since $w \geq 1$ and $\delta < 0$, taking the supremum gives

$$\begin{aligned}\|T(h_1)-T(h_2)\|_{C^0_{\delta-4}} &\leq \Big\{\|h_1-h_2\|_{C^0_\delta}\|h_1\|_{C^4_{\delta-4}}+\|h_2\|_{C^0_\delta}\|h_1-h_2\|_{C^4_{\delta-4}}\Big\}\\ &\leq C(\|h_1\|_{C^4_\delta}+\|h_2\|_{C^4_\delta})\cdot\|h_1-h_2\|_{C^4_\delta},\end{aligned}\tag{3.54}$$

Next, consider the term

$$\rho(h)=g*g^{-4}*Rm_g*h*\nabla^2 h.$$

Taking differences as we did above yields

$$|\rho(h_1)-\rho(h_2)|\leq |Rm_g||h_1-h_2||\nabla^2 h_1|+|Rm_g||h_2||\nabla^2(h_1-h_2)|.$$

Multiplying by the appropriate power of the weight,

$$\begin{aligned}|\rho(h_1)-\rho(h_2)|w^{4-\delta} &\leq \big\{w^2|Rm_g|\big\}\big\{|h_1-h_2|w^{-\delta}\big\}\big\{|\nabla^2 h_1|w^{2-\delta}\big\}w^{\delta}\\ &\quad+\big\{w^2|Rm_g|\big\}\big\{|h_2|w^{-\delta}\big\}\big\{|\nabla^2(h_1-h_2)|w^{2-\delta}\big\}w^{\delta}.\end{aligned}$$

Using (3.29), we arrive at an estimate similar to (3.54).

Finally, let us consider a term in Qh which has a higher order of homogeneity,

$$K(h)=g*r_6(h)*\nabla h*\nabla h*\nabla h*\nabla h.\tag{3.55}$$

Then

$$\begin{aligned}K(h_1)-K(h_2)&=g*[r_6(h_1)-r_6(h_2)]*\nabla h_1*\nabla h_1*\nabla h_1*\nabla h_1\\ &+g*r_6(h_2)\big\{\nabla(h_1-h_2)*\nabla h_1*\nabla h_1*\nabla h_1+\nabla h_2*\nabla(h_1-h_2)*\nabla h_1*\nabla h_1\\ &+\nabla h_2*\nabla h_2*\nabla(h_1-h_2)*\nabla h_1+\nabla h_2*\nabla h_2*\nabla h_2*\nabla(h_1-h_2)\big\}\end{aligned}$$

Multiplying by the weight,

$$\begin{aligned}|K(h_1)-K(h_2)|w^{4-\delta} &\leq C\{|h_1-h_2|w^{-\delta}\}\{|\nabla h_1|w^{1-\delta}\}^4 w^{4\delta}\\ &\quad+C|\nabla(h_1-h_2)|w^{1-\delta}\Big[\{|\nabla h_1|w^{1-\delta}\}^3+\{|\nabla h_2|w^{1-\delta}\}\{|\nabla h_1|w^{1-\delta}\}^2\\ &\quad+\{|\nabla h_1|w^{1-\delta}\}\{|\nabla h_2|w^{1-\delta}\}^2+\{|\nabla h_1|w^{1-\delta}\}^3\Big]w^{3\delta},\end{aligned}$$

which gives an estimate as in (3.54).

Similar arguments (estimating difference quotients) give the Holder estimate in (3.30).

□

Since the operator P_g differs from B^t only by the gauge term, a similar estimate holds for B^t, see the following Proposition. This fact will be used in several places below (e.g., Proposition 10.3).

Proposition 3.8. *Let $(B^t_g)'$ denote the linearization of the B^t tensor:*

$$(B^t_g)'h = \frac{d}{ds}B^t(g+sh)\big|_{s=0}.$$

If we write

$$B^t(g+h) = B^t(g) + (B^t_g)'h + \mathcal{Q}_g(h), \tag{3.56}$$

then under the same assumptions as in (i) of Proposition 3.7, the remainder $\mathcal{Q}$ satisfies the estimate (3.27)*.*

4. Cokernel on a compact manifold

On a compact manifold (Z, g_Z), with basepoint z_0, we define the weight function to be a smooth function satisfying

$$w(z) = \begin{cases} d(z,z_0) & d(z,z_0) < 1/2 \\ 1 & d(z,z_0) \geq 1, \end{cases} \tag{4.1}$$

and $1/2 \leq w(z) \leq 1$ when $1/2 \leq d(z,z_0) \leq 1$.

Theorem 4.1. *Let (Z, g_Z) be either $\mathbb{CP}^2$ with the Fubini-Study metric g_{FS}, or $S^2 \times S^2$ with the product metric $g_{S^2} \times g_{S^2}$. Assume that*

$$t < 0, \tag{4.2}$$

and let $h \in C^{4,\alpha}_{\delta}$ solve the equation

$$S^t(h) = 0 \tag{4.3}$$

for $\delta < 0$ with $|\delta|$ small. If h is toric-invariant and diagonally invariant, then $h = c \cdot g_Z$ for some constant $c \in \mathbb{R}$. Consequently, if h satisfies

$$h = O(|z|^{\delta}) \tag{4.4}$$

as $|z| \to 0$, for $\delta > 0$, then $h \equiv 0$.

Proof. For $t \neq 0$, we define H^1_t to be the kernel of the linearization of P_g:

$$H^1_t = H^1_t(M,g) = \big\{ h \in \overline{S}^2_0(T^*M) \;\big|\; S^t_g h = 0 \big\}, \tag{4.5}$$

where

$$\overline{S}^2_0(T^*M) = \Big\{ h \in C^{4,\alpha}(S^2(T^*M)) \;:\; \int (tr_g\ h)\, dV_g = 0 \Big\}. \tag{4.6}$$

For $t = 0$ (the Bach tensor), we restrict to traceless tensors:

$$H^1_0 = H^1_0(M,g) = \big\{ h \in C^{4,\alpha}(S^2_0(T^*M)) \;\big|\; S^0_g h = 0 \big\}. \tag{4.7}$$

If $H^1_t(M,g) = \{0\}$, we say that (M,g) is *infinitesimally B^t-rigid.* We next quote two crucial rigidity theorems from [GV11] with the following caveat: as pointed out in the proof of Proposition 3.3, a different parametrization τ was used in [GV11]. The relation between τ and t is given by

$$\tau = \frac{t}{2} - \frac{1}{3}. \tag{4.8}$$

The following is then a direct consequence of [GV11, Theorem 7.8]:

Theorem 4.2 ([GV11])**.** *On* $(\mathbb{CP}^2, g_{FS})$, $H^1_t = 0$ *provided that* $t < 1$.

The following is a direct consequence of [GV11, Theorem 7.13]:

Theorem 4.3 ([GV11])**.** *On* $(S^2 \times S^2, g_{S^2 \times S^2})$, $H^1_t = 0$ *provided that* $t < 2/3$ *and* $t \neq -1/3$. *If* $t = -1/3$, *then* H^1_t *is one-dimensional and spanned by the element* $g_1 - g_2$.

If one knows that $h \in C^{4,\alpha}(Z)$, then Theorem 4.1 follows immediately from Theorems 4.2 and 4.3. The only symmetry needed for this part is the diagonal invariance for $t = -1/3$, which rules out the kernel element $g_1 - g_2$. We will next employ the symmetries, in a crucial way, to prove smoothness.

Proposition 4.4. *If* $t \neq 0$, *the indicial roots of* S^t *are contained in* $\mathbb{Z}$.

Proof. To determine the indicial roots of S^t, we need to analyze homogeneous solutions of the equation

$$\begin{aligned} S_0 h \equiv \Delta_0^2 h - 2\big(t + \frac{5}{24}\big)\nabla_0^2(\Delta_0 tr\ h) - 2\big(t + \frac{5}{24}\big)\Delta_0(\delta_0^2 h) g_0 \\ + 2\big(t + \frac{5}{6}\big)\nabla_0^2(\delta_0^2 h) + 2\big(t - \frac{7}{96}\big)\Delta_0^2(tr\ h) g_0 = 0 \end{aligned} \tag{4.9}$$

on Euclidean space $(\mathbb{R}^4 \setminus \{0\}, g_0)$. Assume by contradiction that h solves (4.9) in $\mathbb{R}^4 \setminus \{0\}$, with h corresponding to an indicial root of $u + \sqrt{-1}v \in \mathbb{C} \setminus \{\mathbb{Z}\}$, and $u, v \in \mathbb{R}$. This means that h has components of the form $r^u \cos(vr)$, $r^u \sin(vr)$, or a polynomial in $\log(r)$ times one of these (we say such a solution is homogeneous of degree $u + \sqrt{-1}v$).

Taking the trace of (4.9) gives

$$\Delta_0\big[\Delta_0(tr\ h) - \delta_0^2 h\big] = 0,$$

with $\Delta_0(tr\ h) - \delta_0^2 h$ homogeneous of degree $u - 2 + \sqrt{-1}v$. Since the indicial roots of the Laplacian are $\mathbb{Z} \setminus \{-1\}$, it follows that

$$\Delta_0(tr\ h) - \delta_0^2 h = 0. \tag{4.10}$$

Substituting this into (4.9) implies that

$$\Delta_0^2 h + \frac{5}{4}\nabla_0^2(\Delta_0 tr\ h) - \frac{9}{16}\Delta_0^2(tr\ h) \cdot g_0 = 0.$$

Applying the operator δ_0^2 and using (4.10) we get

$$\Delta_0^2(\delta_0^2 h) = 0,$$

which implies $\Delta_0 tr\ h = \delta_0^2 h = 0$, hence $\Delta_0^2 h \equiv 0$. We note that the indicial roots of Δ_0 on symmetric tensors are the same as those of the Laplacian on functions, which is $\mathbb{Z} \setminus \{-1\}$. Since $u + \sqrt{-1}v$ is not an indicial root of Δ_0^2 on symmetric tensors, we have a contradiction. $\square$

To analyze the indicial root at 0, we first note that any constant tensor on $\mathbb{R}^4$ is a homogeneous degree zero solution, and the dimension of the space of these solutions is 10. We claim that the space of all homogeneous solutions of degree zero is of dimension 20. To see this, choose weight function on $\mathbb{R}^4$ to be given by

$$w(x) = \begin{cases} |x| & |x| \geq 1 \\ 1 & d(x, x_0) < 1. \end{cases} \tag{4.11}$$

With this weight function, for $\delta > 0$ small but nonzero, consider the operator as mapping from

$$S_{g_0}^t : C_\delta^{4,\alpha} \to C_{\delta-4}^{0,\alpha}. \tag{4.12}$$

With obvious notation, the relative index theorem of [LM85] states that

$$Ind(\delta) - Ind(-\delta) = N(0), \tag{4.13}$$

where $N(0)$ is the space of all homogeneous solutions of degree zero on $\mathbb{R}^4 \setminus \{0\}$. We note the important fact that any bounded solution globally defined on $\mathbb{R}^4$ must be constant, the proof is as in [AV12b, Proposition 5.4] (the key being that the flat metric is rigid). This implies that any globally defined decaying solution is trivial, so we have $\dim Ker(-\delta) = 0$. Since the adjoint weight of δ is $-\delta$, (4.13) may then be written as

$$2 \cdot \dim Ker(\delta) = N(0). \tag{4.14}$$

If δ is sufficiently small, it is not an indicial root, so any kernel element defined on all of $\mathbb{R}^4$ satisfying $h = O(|x|^\delta)$ as $|x| \to \infty$ is constant. Therefore $\dim Ker(\delta) = 10$, which implies that $N(0) = 20$.

The only symmetric constant tensors invariant under the standard diagonal torus action are multiples of the identity matrix, or multiples of the matrix

$$\begin{pmatrix} I_2 & 0 \\ 0 & -I_2 \end{pmatrix}, \tag{4.15}$$

where I_2 is the 2×2 identity matrix. It is easy to see that this element is not invariant under the diagonal symmetry. Consequently, there are only 2 invariant degree zero solutions on $\mathbb{R}^4 \setminus \{0\}$: the identity matrix, and another solution with log-type growth (we will not need the explicit formula). Another application of the relative index theorem applied to the compact manifold (details are similar to above) shows that, since $c \cdot g$ extends to a global solution, the log-type solution does not extend to a global solution on $Z \setminus \{z_0\}$.

To finish the proof, if $h \in C^{4,\alpha}_{\delta}$ is a solution on $Z \backslash \{z_0\}$ for $\delta < 0$ with $|\delta|$ sufficiently small which is invariant under the group action, then there is an expansion

$$h = c \cdot g + O(|z|^{\epsilon}) \tag{4.16}$$

for some constant $c \in \mathbb{R}$ and $\epsilon > 0$ as $|z| \to 0$. Since the leading term is a global solution, we then have that $\tilde{h} = h - c \cdot g$ is solution on $Z \backslash \{z_0\}$ satisfying $\tilde{h} = O(|z|^{\epsilon})$ as $|z| \to 0$. A standard integration-by-parts argument shows that $\tilde{h}$ extends to a weak solution on all of Z, and is therefore smooth by elliptic regularity. By the above, $\tilde{h} \equiv 0$. $\square$

5. Cokernel on an asymptotically flat manifold

Let (N, g) be the Green's function metric of a compact manifold (Y, g_Y) with positive scalar curvature: more precisely,

$$\begin{aligned} N &= Y \setminus \{y_0\},\ y_0 \in Y; \\ g &= G^2 g_Y, \end{aligned} \tag{5.1}$$

where G is the Green's function of the conformal Laplacian with pole at $y_0 \in Y$. Assume (Y, g_Y) is Bach-flat and infinitesimally Bach-rigid, that is, $H^1_0(Y, g_Y) = \{0\}$.

Let $\{x^i\}$ denote an inverted normal coordinate system, and choose weight function $w = w(x)$ to be given by

$$w(x) = \begin{cases} |x| & |x| \geq R_0 \\ 1 & d(x, x_0) < 1, \end{cases} \tag{5.2}$$

where R_0 is large, and $x_0 \in N$ is a basepoint.

Theorem 5.1. *Assume $\delta < 0$ with $|\delta|$ small, and let $h \in C^{4,\alpha}_{\delta}$ solve the equation*

$$S^t(h) = B'(h) + tC'(h) + \mathcal{K}\delta\mathcal{K}\delta(\overset{\circ}{h}) = 0, \tag{5.3}$$

where $t \neq 0$.

Then

$$h = \mathcal{K}\omega_1 + f \cdot g_N, \tag{5.4}$$

where ω_1 and f satisfy

$$\begin{aligned} \square\omega_1 &= 0, \\ \Delta f &= -\frac{1}{3}\langle Ric, \mathcal{K}\omega_1 \rangle, \end{aligned} \tag{5.5}$$

where $\square = \delta\mathcal{K}$.

Furthermore, suppose (N, g) is either the Burns metric or the Green's function metric of the product metric on $S^2 \times S^2$. If h is toric invariant and diagonally *invariant and* $\delta > 0$*, then* ω *and* f *can also be chosen to be toric invariant and diagonally invariant, with*

$$\omega = c \cdot x^i dx^i + O(|x|^{-1+\epsilon}), \tag{5.6}$$

$$f(x) = c_0 + \frac{c'}{|x|^2} + O'(|x|^{-4+\epsilon}), \tag{5.7}$$

where $c_0, c, c' \in \mathbb{R}$ *are constants, as* $r \to \infty$*, for any* $\epsilon > 0$*.*

The remainder of this section will be devoted to the proof of Theorem 5.1. Since the Bach tensor is conformally invariant it follows that (N, g) is also Bach-flat. Also, since (N, g) is scalar flat it is also B^t-flat, for any value of t. We also note that h is smooth since S^t is elliptic.

The splitting in (5.4) reflects the fact that each term in the linearization must vanish:

Proposition 5.2. *Each term in (5.3) vanishes; i.e.,*

$$\begin{aligned} B'(h) &= 0, \\ C'(h) &= 0, \\ \mathcal{K}\delta\mathcal{K}\delta(\mathring{h}) &= 0. \end{aligned} \tag{5.8}$$

Furthermore,

$$\delta(\mathring{h}) = 0. \tag{5.9}$$

Proof. Since (N, g) is B^t-flat, if we linearize the identity

$$\delta B^t = 0$$

at g we find

$$\delta[(B^t)'h] + (\delta'_h)B^t = 0 \ \Rightarrow \ \delta[(B^t)'h] = 0.$$

Therefore, taking the divergence of both sides of (5.3) gives

$$\Box^2 \delta(\mathring{h}) = 0. \tag{5.10}$$

Proposition 5.3. *There are no decaying elements in the kernel of* $\Box$*.*

Proof. To see this, we note the formula

$$\Box\omega = \frac{3}{2} d\delta\omega + \delta d\omega + 2Ric(\omega, \cdot). \tag{5.11}$$

Since the Ricci tensor decays, to determine the indicial roots of $\Box$, we need to analyze homogeneous solutions of the operator

$$\Box\omega = \frac{3}{2} d\delta\omega + \delta d\omega \tag{5.12}$$

on Euclidean space $(\mathbb{R}^4, g_0)$. We claim that the indicial roots are contained in $\mathbb{Z}\backslash\{-1\}$. To prove this, assume by contradiction that ω solves $\Box\omega = 0$ in $\mathbb{R}^4 \setminus \{0\}$, with ω corresponding to an indicial root of $u + \sqrt{-1}v \in \mathbb{C} \setminus \{\mathbb{Z} \setminus \{-1\}\}$, and $u, v \in \mathbb{R}$. This means that ω has components of the form $r^u \cos(vr)$ or $r^u \sin(vr), r^u \log(r)$ (similarly to above, we say such a solution is homogeneous of degree $u + \sqrt{-1}v$). Applying d to (5.12) yields that

$$d\delta d\omega = (d\delta + \delta d)d\omega = -\Delta_H d\omega. \tag{5.13}$$

We note that the indicial roots of Δ_H are exactly $\mathbb{Z} \setminus \{-1\}$ (this is easily seen since the leading term is the rough Laplacian, so the indicial roots are the same as for the Laplacian on functions). Since $d\omega$ is homogeneous of degree $u - 1 + \sqrt{-1}v$, which is not an indicial root of Δ_H, we conclude that $d\omega = 0$. A similiar argument shows that $\delta\omega = 0$. Since both $d\omega = 0$ and $\delta\omega = 0$, we have that $\Delta_H(\omega) = 0$, which is a contradiction since $u + \sqrt{-1}v$ was chosen to not be an indicial root of Δ_H.

Consequently, by standard weighted space theory, any decaying solution of $\Box\xi = 0$ on an AF space must satisfy $\xi = O(r^{-2})$ as $r \to \infty$ [Bar86]. An elementary integration by parts argument then shows that $\mathcal{K}\xi = 0$. As there are no decaying conformal Killing fields on an AF space, we conclude that $\xi = 0$. $\Box$

Remark 5.4. By a separation of variables argument as in [AV12b, Section 4.1], it is straightforward to show that the indicial roots of $\Box$ are in fact exactly $\mathbb{Z} \setminus \{-1\}$, although we will not need this fact.

By this proposition and (5.10),

$$\Box\delta(\overset{\circ}{h}) = 0.$$

Applying the result once again gives (5.9):

$$\delta\overset{\circ}{h} = 0. \tag{5.14}$$

In particular,

$$\mathcal{K}\delta\mathcal{K}\delta(\overset{\circ}{h}) = 0$$

and consequently

$$B'(h) + tC'(h) = 0. \tag{5.15}$$

If we linearize the trace-free property of the Bach tensor at g it follows that

$$trB'(h) = 0.$$

Therefore, taking the trace of (5.15) gives

$$\begin{aligned} 0 &= tr\ B'(h) + t\ tr\ C'(h) \\ &= t\ tr\ C'(h). \end{aligned} \tag{5.16}$$

Lemma 5.5. *If (X^4, g) is either scalar-flat or Einstein, then*

$$\begin{aligned} tr\ C'(h) &= -6\Delta R'(h) \\ &= -6\Delta[-\Delta(tr\ h) + \delta^2 h - \langle Ric, h\rangle], \end{aligned} \tag{5.17}$$

where R' denotes the linearization of the scalar curvature.

Proof. Since $C = 0$ for scalar-flat or Einstein metrics, we have

$$\begin{aligned}(tr\ C)' &= (tr\ )'C + tr(C') \\ &= tr(C').\end{aligned}$$

Also, $tr\ C = -6\Delta R$, so

$$tr(C') = (tr\ C)' = -6(\Delta)'R - 6\Delta R',$$

and since R is constant we get

$$tr(C') = -6\Delta R',$$

as claimed. ☐

In view of (5.16) and the preceding lemma we have

$$\Delta R'(h) = \Delta[-\Delta(tr\ h) + \delta^2 h - \langle Ric, h\rangle] = 0.$$

Since

$$|\nabla^2 h| = O(|x|^{\delta-2}), \quad \langle Ric, h\rangle = O(|x|^{\delta-4}),$$

it follows that $R'(h)$ is a decaying harmonic function. Therefore,

$$R'(h) = -\Delta(tr\ h) + \delta^2 h - \langle Ric, h\rangle = 0. \tag{5.18}$$

Recall C is given by

$$C = 2\nabla^2 R - 2(\Delta R)g - 2R\big(Ric - \frac{1}{4}Rg\big). \tag{5.19}$$

Linearizing this at g (which is scalar-flat) gives

$$C'(h) = 2\nabla^2 R'(h) - 2\Delta R'(h)g - 2R'(h)\cdot Ric.$$

From (5.18), it follows that $C'(h) = 0$, which completes the proof of Proposition 5.2. ☐

Write

$$h = \overset{\circ}{h} + fg, \tag{5.20}$$

where $f = (tr\ h)/4$. The conformal invariance of the Bach tensor leads to the formula

$$B'(\phi g) = -2\phi B$$

for any function ϕ. Since g is Bach-flat this implies

$$\begin{aligned}0 &= B'(\overset{\circ}{h} + fg) \\ &= B'(\overset{\circ}{h}).\end{aligned}$$

It follows from [AV12b, Proposition 2.1] that any decaying, transverse-traceless element in the kernel of B' must decay quadratically, hence

$$|\overset{\circ}{h}| = O(|x|^{-2}), \tag{5.21}$$

as $|x| \to \infty$.

Conformal invariance of the Bach tensor also implies the invariance of its linearization:

$$0 = B'_g(\overset{\circ}{h}) = B'_{G^2 g_0}(\overset{\circ}{h}) = G^{-2} B'_{g_0}(G^{-2}\overset{\circ}{h}). \tag{5.22}$$

Denote

$$\tilde{h} = G^{-2}\overset{\circ}{h}. \tag{5.23}$$

Then $\tilde{h} \in C^{\infty}(Y \setminus \{y_0\})$, and

$$B'_{g_0}\tilde{h} = 0. \tag{5.24}$$

In addition, since $\overset{\circ}{h}$ decays quadratically at infinity, $\tilde{h}$ vanishes quadratically at y_0. To see this, first note that

$$\begin{aligned} |\tilde{h}|^2_{g_0} &= (g_0)^{ik}(g_0)^{k\ell}\tilde{h}_{ij}\tilde{h}_{k\ell} \\ &= G^4 g^{ik} g^{j\ell}\big(G^{-2}\overset{\circ}{h}_{ij}\big)\big(G^{-2}\overset{\circ}{h}_{k\ell}\big) \\ &= |\overset{\circ}{h}|^2_g. \end{aligned} \tag{5.25}$$

We note the relation between $r = |x|$ and $\rho = |y|$:

$$r \sim \rho^{-1},$$

so that (5.21) and (5.25) together imply

$$|\tilde{h}|_{g_0} = O(\rho^2), \tag{5.26}$$

as $\rho \to 0$. In particular, $\tilde{h} \in C^{1,\alpha}(Y)$.

We now use the standard splitting of a trace-free symmetric tensor into the image of the conformal Killing operator and the space of transverse-traceless tensors. More precisely, we first solve

$$\Box_{g_0}\omega_0 = \delta_{g_0}\tilde{h} \tag{5.27}$$

with $\omega_0 \in C^{2,\alpha}(Y)$. Since $\Box$ is self-adjoint with kernel given by the space of conformal Killing forms $\mathcal{C}(Y, g_0)$, this equation is solvable whenever the right-hand side is orthogonal to $\mathcal{C}(Y, g_0)$. However, if $\eta \in \mathcal{C}(Y, g_0)$, then

$$\int \langle \delta_{g_0}\tilde{h}, \eta\rangle \, dV_0 = -\frac{1}{2}\int \langle \tilde{h}, \mathcal{K}_{g_0}\eta\rangle \, dV_0 = 0.$$

It follows that (5.27) is always solvable, although the solution ω_0 is only unique up to the space of conformal Killing fields. This fact will actually be crucial when we impose toric and diagonal invariance, in which case we will need to solve (5.27) equivariantly and study the space of invariant forms (see the end of this section).

Let

$$h_0 = \tilde{h} - \mathcal{K}_{g_0}\omega_0. \tag{5.28}$$

Then $h_0 \in C^{1,\alpha}(Y)$, and is smooth away from y_0. By (5.27), h_0 is transverse-traceless, and on $Y \setminus \{y_0\}$

$$\begin{aligned} B'_{g_0}(h_0) &= B'_{g_0}(\tilde{h} - \mathcal{K}_{g_0}\omega_0) \\ &= B'_{g_0}(\tilde{h}) \ \ (\text{since Im } \mathcal{K} \subset \text{Ker} B') \\ &= 0. \end{aligned}$$

A standard integration by parts argument shows that h_0 is a global weak solution of $B'_{g_0} h_0 = 0$, and from elliptic theory it follows that h_0 is smooth on Y. Since (Y, g_0) is assumed to be infinitesimally Bach-rigid, $h_0 = 0$, and we conclude that

$$\tilde{h} = \mathcal{K}_{g_0}\omega_0. \tag{5.29}$$

By conformal invariance of the conformal Killing operator,

$$\mathcal{K}_{g_0}\omega_0 = G^{-2}\mathcal{K}_g[G^2\omega_0]. \tag{5.30}$$

Hence,

$$G^{-2}\overset{\circ}{h} = \tilde{h} = G^{-2}\mathcal{K}_g[G^2\omega_0],$$

which implies

$$\overset{\circ}{h} = \mathcal{K}_g[G^2\omega_0]. \tag{5.31}$$

Also, by (5.14), $\omega_1 = G^2\omega_0$ satisfies

$$0 = \delta\overset{\circ}{h} = \Box\omega_1, \tag{5.32}$$

which gives the first equation in (5.5). To prove the second equation, use the splitting $h = \overset{\circ}{h} + fg$ in (5.18); this gives

$$-3\Delta f - \langle Ric, \overset{\circ}{h}\rangle = 0 \tag{5.33}$$

(note we have used the scalar-flat condition again).

Up to this point we have not used the invariance of h. In general, the form ω_1 can grow quadratically on N; however, using invariance we can choose a solution ω_0 of (5.27) so that the resulting form ω_1 has linear growth on N, with highest order given by (5.6). To see this, we argue as follows. Since $\omega_0 \in C^{2,\alpha}(Y)$, it admits an expansion

$$\omega_0 = \omega^{(0)} + \omega^{(1)} + \omega^{(2)} + O(\rho^{2+\alpha}), \tag{5.34}$$

as $\rho \to 0$, where

$$\omega^{(0)} = \omega^{(0)}_i dy^i \tag{5.35}$$

$$\omega^{(1)} = \omega^{(1)}_{ij} y^i dy^j \tag{5.36}$$

$$\omega^{(2)} = \omega^{(2)}_{ijk} y^i y^j dy^k, \tag{5.37}$$

where the $\{y\}$-coordinates are local normal coordinates near y_0 with torus action

$$(y_1, y_2, y_3, y_4) \mapsto \big(e^{\sqrt{-1}\theta_1}(y_1 + \sqrt{-1}y_2), e^{\sqrt{-1}\theta_2}(y_3 + \sqrt{-1}y_4)\big). \tag{5.38}$$

Denote $\rho_1^2 = y_1^2 + y_2^2$, $\rho_2^2 = y_3^2 + y_4^2$, and $\rho^2 = \rho_1^2 + \rho_2^2$, and let θ_1, θ_2 denote the corresponding angular coordinates. Since the group is compact, we can average over the group to find a solution of (5.27) which is also invariant under the group action (5.38), as well as the diagonal symmetry. It is elementary to see that there is no form with constant coefficients which is invariant under the torus action (5.38). The only toric-invariant 1-forms with linear coefficients are

$$c_1 d\rho_1 + c_2 \rho_1 d\theta_1 + c_3 d\rho_2 + c_4 \rho_2 d\theta_2. \tag{5.39}$$

The forms $\rho_1 d\theta_1$ and $\rho_2 d\theta_2$ extend to global Killing forms, so we may assume that $c_2 = c_4 = 0$. Invariance under the diagonal symmetry implies that $c_1 = c_3$, so we have that

$$\omega_0 = c \cdot \rho d\rho + \omega^{(2)} + O(\rho^{2+\alpha}), \tag{5.40}$$

for some constant $c \in \mathbb{R}$. This implies the expansion

$$\omega_1 = c \cdot x^i dx^i + O(|x|^{-\alpha}), \tag{5.41}$$

as $|x| \to \infty$. Averaging over the group, we may assume ω_1 is also invariant under the group action.

To obtain the expansion for f, extend the function $|x|^{-2}$ to all of N by a cutoff function (which we supress). It is not hard to see that $\Delta(|x|^{-2}) = O(|x|^{-6})$ as $|x| \to \infty$, and there exists a constant $c' \in \mathbb{R}$ so that

$$\int_N \Big(-\frac{1}{3}\langle Ric, \mathcal{K}\omega_1\rangle - c'\Delta|x|^{-2}\Big) dV = 0. \tag{5.42}$$

Next, consider $\Delta : C^{2,\alpha}_{-4+\epsilon}(N) \to C^{0,\alpha}_{-6+\epsilon}(N)$. The adjoint weight is $2-\epsilon$, so from toric invariance, the kernel of the adjoint contains only constants. We may then solve the equation

$$\Delta \tilde{f} = -\frac{1}{3}\langle Ric, \mathcal{K}\omega_1\rangle - c'\Delta(|x|^{-2}) \tag{5.43}$$

with $\tilde{f} \in C^{2,\alpha}_{-4+\epsilon}$. Equivalently,

$$\Delta(\tilde{f} + c'|x|^{-2}) = -\frac{1}{3}\langle Ric, \mathcal{K}\omega_1\rangle. \tag{5.44}$$

Since there are no decaying harmonic functions, we must have

$$f = \tilde{f} + c'|x|^{-2} \tag{5.45}$$

with $\tilde{f} = O(|x|^{-4+\epsilon})$ as $|x| \to \infty$ for any $\epsilon > 0$. Again, averaging over the group, we may assume that f is invariant under the group action.

Finally, we consider the case that $\delta > 0$. Using the same argument as in the proof of Theorem 4.1 involving the relative index theorem, the toric and diagonal symmetries imply that the only possible leading terms are $c \cdot g_N$ and a log-type solution. Since $c \cdot g_N$ extends to a global solution, again the relative index theorem implies that the log-type solution does not occur. Consequently, after subtracting a multiple of the metric, the solution is decaying, and (5.7) follows from the previous expansion.

6. Asymptotics of the cokernel

Denote the (normalized) cokernel element described in Theorem 5.1 by

$$o_1 = \kappa + fg, \tag{6.1}$$

where

$$\kappa = \mathcal{K}[\omega_1], \tag{6.2}$$

with

$$(\omega_1)_i = x^i + O(1). \tag{6.3}$$

This section will be devoted to proving the following

Theorem 6.1. *The tracefree part of AF-cokernel element o_1 satisfies*

$$\kappa_{ij} = \frac{2}{3} W_{ikj\ell}(y_0) \frac{x^k x^\ell}{|x|^4} + O(|x|^{-4+\epsilon}) \tag{6.4}$$

as $|x| \to \infty$, for any $\epsilon > 0$.

Recall in inverted normal coordinates at the point y_0, the AF metric has the expansion

$$g_{ij} = \delta_{ij} - \frac{1}{3} R_{ikj\ell}(y_0) \frac{x^k x^\ell}{|x|^4} + \frac{2A}{|x|^2}\delta_{ij} + O(|x|^{-3}). \tag{6.5}$$

In the following, we will need to have expansions for the Christoffel symbols:

Lemma 6.2. *In inverted normal coordinates,*

$$\begin{aligned} \Gamma_{ij}^k = &- \frac{1}{3} R_{i\alpha kj}(y_0) \frac{x^\alpha}{|x|^4} - \frac{1}{3} R_{j\alpha ki}(y_0) \frac{x^\alpha}{|x|^4} \\ &+ \frac{2}{3} R_{i\alpha k\beta}(y_0) \frac{x^j x^\alpha x^\beta}{|x|^6} + \frac{2}{3} R_{j\alpha k\beta}(y_0) \frac{x^i x^\alpha x^\beta}{|x|^6} - \frac{2}{3} R_{i\alpha j\beta}(y_0) \frac{x^k x^\alpha x^\beta}{|x|^6} \\ &- 2A \frac{x^i \delta_{jk}}{|x|^4} - 2A \frac{x^j \delta_{ik}}{|x|^4} + 2A \frac{x^k \delta_{ij}}{|x|^4} + O(|x|^{-4}). \end{aligned} \tag{6.6}$$

Proof. Recall

$$\Gamma_{ij}^k = \frac{1}{2} g^{km} \big(\partial_i g_{jm} + \partial_j g_{im} - \partial_m g_{ij} \big).$$

By (6.5),

$$\begin{aligned} \partial_i g_{jm} &= \partial_i \Big\{ - \frac{1}{3} R_{j\alpha m\beta}(y_0) \frac{x^\alpha x^\beta}{|x|^4} + \frac{2A}{|x|^2} \delta_{jm} \Big\} + \cdots \\ &= -\frac{1}{3} R_{jim\beta}(y_0) \frac{x^\beta}{|x|^4} - \frac{1}{3} R_{j\alpha mi}(y_0) \frac{x^\alpha}{|x|^4} + \frac{4}{3} R_{j\alpha m\beta}(y_0) \frac{x^i x^\alpha x^\beta}{|x|^6} \\ &\quad - 4A \frac{x^i \delta_{jm}}{|x|^4} + \cdots \end{aligned}$$

Therefore, after combining terms and rearranging,

$$\begin{aligned}\partial_i g_{jm} + \partial_j g_{im} - \partial_m g_{ij} = &-\frac{2}{3}R_{j\alpha mi}(y_0)\frac{x^\alpha}{|x|^4} - \frac{2}{3}R_{i\alpha mj}(y_0)\frac{x^\alpha}{|x|^4}\\ &+\frac{4}{3}R_{j\alpha m\beta}(y_0)\frac{x^i x^\alpha x^\beta}{|x|^6} + \frac{4}{3}R_{i\alpha m\beta}(y_0)\frac{x^j x^\alpha x^\beta}{|x|^6} - \frac{4}{3}R_{j\alpha i\beta}(y_0)\frac{x^m x^\alpha x^\beta}{|x|^6}\\ &-4A\frac{x^i\delta_{jm}}{|x|^4} - 4A\frac{x^j\delta_{im}}{|x|^4} + 4A\frac{x^m\delta_{ij}}{|x|^4} + \cdots\end{aligned}$$

It follows from (6.5) that the inverse matrix g^{km} is given by

$$g^{km} = \delta_{km} + \frac{1}{3}R_{k\alpha m\beta}(y_0)\frac{x^\alpha x^\beta}{|x|^4} - \frac{2A}{|x|^2}\delta_{km} + \cdots \tag{6.7}$$

Consequently,

$$\begin{aligned}\Gamma^k_{ij} = &-\frac{1}{3}R_{j\alpha ki}(y_0)\frac{x^\alpha}{|x|^4} - \frac{1}{3}R_{i\alpha kj}(y_0)\frac{x^\alpha}{|x|^4}\\ &+\frac{2}{3}R_{j\alpha k\beta}(y_0)\frac{x^j x^\alpha x^\beta}{|x|^6} + \frac{2}{3}R_{i\alpha k\beta}(y_0)\frac{x^j x^\alpha x^\beta}{|x|^6} - \frac{2}{3}R_{j\alpha i\beta}(y_0)\frac{x^k x^\alpha x^\beta}{|x|^6}\\ &-2A\frac{x^i\delta_{jk}}{|x|^4} - 2A\frac{x^j\delta_{ik}}{|x|^4} + 2A\frac{x^k\delta_{ij}}{|x|^4} + O(|x|^{-4}),\end{aligned}\tag{6.8}$$

which is the same as (6.6) after rearranging and re-indexing. □

Next, we consider the form $\omega = \omega_j dx^j$ with

$$\omega_j = x^j + b_{jk}\frac{x^k}{|x|^2}, \tag{6.9}$$

where

$$b_{ij} = -\frac{1}{3}S_{ij}(y_0) + 2A\delta_{ij}, \tag{6.10}$$

where

$$S_{ij}(y_0) = \frac{1}{2}\Big(R_{ij}(y_0) - \frac{1}{6}R(y_0)\delta_{ij}\Big) \tag{6.11}$$

is the Schouten tensor. We extend ω to be a globally defined form on all of N by cutting it off at some finite distance from the basepoint. Since this cutoff will not matter in the following, we will suppress it from the following computations.

Lemma 6.3. *In inverted normal coordinates,*

$$\begin{aligned}\mathcal{K}[\omega] = &\,2\frac{b_{ij}}{|x|^2} - 2\frac{b_{jk}x^k x^i}{|x|^4} - 2\frac{b_{ik}x^k x^j}{|x|^4} - \frac{1}{2}\frac{b_{kk}}{|x|^4}\delta_{ij} + \frac{b_{k\ell}x^k x^\ell}{|x|^4}\delta_{ij}\\ &+\frac{2}{3}R_{ikj\ell}(y_0)\frac{x^k x^\ell}{|x|^4} - \frac{1}{6}R_{k\ell}(y_0)\frac{x^k x^\ell}{|x|^4}\delta_{ij}\\ &+\frac{2A}{|x|^4}\big[4x^i x^j - |x|^2\delta_{ij}\big] + O(|x|^{-4}),\end{aligned}\tag{6.12}$$

as $|x| \to \infty$.

Proof. We begin by noting

$$\nabla_i \omega_j = \partial_i \omega_j - \Gamma_{ij}^k \omega_k.$$

Using (6.9),

$$\partial_i \omega_j = \delta_{ij} + \frac{b_{ji}}{|x|^2} - 2\frac{b_{jk}x^k x^i}{|x|^4} + \cdots , \tag{6.13}$$

while

$$\begin{aligned} \Gamma_{ij}^k \omega_k = & -\frac{1}{3} R_{i\alpha kj}(y_0)\frac{x^\alpha x^k}{|x|^4} - \frac{1}{3} R_{j\alpha ki}(y_0)\frac{x^\alpha x^k}{|x|^4} \\ & + \frac{2}{3} R_{i\alpha k\beta}(y_0)\frac{x^j x^k x^\alpha x^\beta}{|x|^4} + \frac{2}{3} R_{j\alpha k\beta}(y_0)\frac{x^j x^k x^\alpha x^\beta}{|x|^4} - \frac{2}{3} R_{i\alpha j\beta}(y_0)\frac{x^\alpha x^\beta}{|x|^2} \\ & - 4A\frac{x^i x^j}{|x|^4} + 2A\frac{\delta_{ij}}{|x|^2} + O(|x|^{-3}). \end{aligned} \tag{6.14}$$

By the symmetries of the curvature tensor, the third and fourth terms above obviously vanish. If we re-index in the first two terms, $k \leftrightarrow \beta$, then we can rewrite them as

$$\begin{aligned} -\frac{1}{3} R_{i\alpha kj}(y_0)\frac{x^\alpha x^k}{|x|^4} - \frac{1}{3} R_{j\alpha ki}(y_0)\frac{x^\alpha x^k}{|x|^4} &= -\frac{1}{3} R_{i\alpha\beta j}(y_0)\frac{x^\alpha x^\beta}{|x|^4} - \frac{1}{3} R_{j\alpha\beta i}(y_0)\frac{x^\alpha x^\beta}{|x|^4} \\ &= \frac{1}{3} R_{i\alpha j\beta}(y_0)\frac{x^\alpha x^\beta}{|x|^4} + \frac{1}{3} R_{i\beta j\alpha}(y_0)\frac{x^\alpha x^\beta}{|x|^4} \\ &= \frac{2}{3} R_{i\alpha j\beta}(y_0)\frac{x^\alpha x^\beta}{|x|^4}. \end{aligned} \tag{6.15}$$

Substituting this back into (6.14), we find that

$$\Gamma_{ij}^k \omega_k = -4A\frac{x^i x^j}{|x|^4} + 2A\frac{\delta_{ij}}{|x|^2} + O(|x|^{-3}).$$

Therefore,

$$\nabla_i \omega_j = \delta_{ij} + \frac{b_{ji}}{|x|^2} - 2\frac{b_{jk}x^k x^i}{|x|^4} + 4A\frac{x^i x^j}{|x|^4} - 2A\frac{\delta_{ij}}{|x|^2} + O(|x|^{-3}). \tag{6.16}$$

The divergence of ω is

$$\begin{aligned} \delta\omega &= g^{ij}\nabla_i \omega_j \\ &= \big\{ \delta_{ij} + \frac{1}{3} R_{i\alpha j\beta}(y_0)\frac{x^\alpha x^\beta}{|x|^4} - \frac{2A}{|x|^2}\delta_{ij} + \cdots \big\} \\ & \qquad \times \big\{ \delta_{ij} + \frac{b_{ji}}{|x|^2} - 2\frac{b_{jk}x^k x^i}{|x|^4} - 2\frac{b_{jk}x^k x^i}{|x|^4} + 4A\frac{x^i x^j}{|x|^4} - 2A\frac{\delta_{ij}}{|x|^2} + \cdots \big\} \\ &= 4 + \frac{b_{kk}}{|x|^2} - 2\frac{b_{k\ell}x^k x^\ell}{|x|^4} + \frac{1}{3} R_{k\ell}(y_0)\frac{x^k x^\ell}{|x|^4} - \frac{12A}{|x|^2} + \cdots . \end{aligned} \tag{6.17}$$

Hence,

$$
\begin{aligned}
(\delta\omega)g_{ij} &= \big\{4 + \frac{b_{kk}}{|x|^2} - 2\frac{b_{k\ell}x^k x^\ell}{|x|^4} + \frac{1}{3}R_{k\ell}(y_0)\frac{x^k x^\ell}{|x|^4} - \frac{12A}{|x|^2} + \cdots\big\} \\
&\quad \times \big\{\delta_{ij} - \frac{1}{3}R_{ikj\ell}(y_0)\frac{x^k x^\ell}{|x|^4} + \frac{2A}{|x|^2}\delta_{ij} + \cdots\big\} \\
&= 4\delta_{ij} + \frac{b_{kk}}{|x|^2}\delta_{ij} - 2\frac{b_{k\ell}x^k x^\ell}{|x|^4}\delta_{ij} - \frac{4}{3}R_{ikj\ell}(y_0)\frac{x^k x^\ell}{|x|^4} + \frac{1}{3}R_{k\ell}(y_0)\frac{x^k x^\ell}{|x|^4}\delta_{ij} \\
&\quad - \frac{4A}{|x|^2}\delta_{ij} + \cdots
\end{aligned}
\tag{6.18}
$$

Finally, combining (6.16) and (6.18) we get

$$
\begin{aligned}
\mathcal{K}[\omega]_{ij} &= \nabla_i\omega_j + \nabla_j\omega_i - \frac{1}{2}(\delta\omega)g_{ij} \\
&= 2\frac{b_{ij}}{|x|^2} - 2\frac{b_{jk}x^k x^i}{|x|^4} - 2\frac{b_{ik}x^k x^j}{|x|^4} - \frac{1}{2}\frac{b_{kk}}{|x|^4}\delta_{ij} + \frac{b_{k\ell}x^k x^\ell}{|x|^4}\delta_{ij} \\
&\quad + \frac{2}{3}R_{ikj\ell}(y_0)\frac{x^k x^\ell}{|x|^4} - \frac{1}{6}R_{k\ell}(y_0)\frac{x^k x^\ell}{|x|^4}\delta_{ij} + 8A\frac{x^i x^j}{|x|^4} - 2A\frac{\delta_{ij}}{|x|^2} + \cdots ,
\end{aligned}
\tag{6.19}
$$

which completes the proof. ☐

This implies the following decay rate for $\Box\omega$:

Lemma 6.4. *In inverted normal coordinates,*

$$\Box\omega = O(|x|^{-4}), \tag{6.20}$$

as $|x| \to \infty$.

Proof. Recall that $\Box$ is given by

$$\Box\omega_j = \big(\delta\mathcal{K}[\omega]\big)_j = g^{ik}\nabla_k\mathcal{K}[\omega]_{ij} = g^{ik}\partial_k\mathcal{K}[\omega]_{ij} + \mathcal{K}[\omega] * \Gamma.$$

Note that

$$(\mathcal{K}[\omega]) * \Gamma \sim |x|^{-5},$$

so it is much lower order than the derivative term. Also,

$$
\begin{aligned}
g^{ik}\partial_k\mathcal{K}[\omega]_{ij} &= \big(\delta_{ik} + O(|x|^{-2})\big)\partial_k\mathcal{K}_{ij}[\omega] \\
&= \partial_i\mathcal{K}[\omega]_{ij} + \text{(lower)}.
\end{aligned}
$$

Consequently, we obtain

$$
\begin{aligned}
(\Box\omega)_j &= -2\frac{b_{ij}x^i}{|x|^4} - 4\frac{b_{jk}x^k}{|x|^4} - \frac{b_{kk}}{|x|^4}x^j + 4\frac{b_{k\ell}x^k x^\ell x^j}{|x|^6} \\
&\quad + \frac{2}{3}R_{k\ell}(y_0)\frac{x^k x^\ell x^j}{|x|^6} - R_{jk}(y_0)\frac{x^k}{|x|^4} + 12A\frac{x^j}{|x|^4} + O(|x|^{-4})
\end{aligned}
\tag{6.21}
$$

as $|x| \to \infty$. Substituting (6.10) into this completes the proof. ☐

Consider $\Box : C^{2,\alpha}_{-2+\epsilon}(N) \to C^{0,\alpha}_{-4+\epsilon}(N)$, with $\epsilon > 0$ small, and consider the equation

$$\Box(\omega') = \Box\omega \tag{6.22}$$

The cokernel of this operator has domain weight $-\epsilon$, so from Proposition 5.3, there is no cokernel. Consequently, (6.22) has a solution $\omega' \in C^{2,\alpha}_{-2+\epsilon}$. The form $\tilde{\omega} = \omega - \omega'$ is then a solution of $\Box\tilde{\omega} = 0$ with expansion

$$\tilde{\omega}_j = x^j + b_{jk}\frac{x^k}{|x|^2} + O(|x|^{-2+\epsilon}), \tag{6.23}$$

for any $\epsilon > 0$. Since $\tilde{\omega}$ and ω_1 have the same leading term, and their difference is decaying, we must have $\tilde{\omega} = \omega_1$, so of course ω_1 admits the same expansion.

Proof of Theorem 6.1. Substituting (6.10) into (6.12) and using the decomposition of the curvature tensor into Weyl and Schouten parts gives

$$\begin{aligned}
\mathcal{K}[\omega]_{ij} &= 2\big\{-\frac{1}{3}S_{ij}(y_0) + 2A\delta_{ij}\big\}\frac{1}{|x|^2} - 2\big\{-\frac{1}{3}S_{jk}(y_0) + 2A\delta_{jk}\big\}\frac{x^i x^k}{|x|^4}\\
&\quad - 2\big\{-\frac{1}{3}S_{ik}(y_0) + 2A\delta_{ik}\big\}\frac{x^j x^k}{|x|^4} - \frac{1}{2}\big\{-\frac{1}{18}R(y_0) + 8A\big\}\frac{\delta_{ij}}{|x|^2}\\
&\quad + \big\{-\frac{1}{3}S_{k\ell}(y_0) + 2A\delta_{k\ell}\big\}\frac{x^k x^\ell}{|x|^4}\delta_{ij} + \frac{2}{3}\big\{W_{ikj\ell}(y_0) + (\delta_{ij}S_{k\ell}(y_0) - \delta_{i\ell}S_{jk}(y_0)\\
&\quad - \delta_{jk}S_{i\ell}(y_0) + \delta_{k\ell}S_{ij}(y_0))\big\}\frac{x^k x^\ell}{|x|^4} + 8A\frac{x^i x^j}{|x|^4} - \frac{1}{6}R_{k\ell}(y_0)\frac{x^k x^\ell}{|x|^4}\delta_{ij} - 2A\frac{\delta_{ij}}{|x|^2} + \cdots\\
&= \frac{2}{3}W_{ikj\ell}(y_0)\frac{x^k x^\ell}{|x|^4} + \cdots
\end{aligned}$$

□

7. Some auxiliary linear equations

In this section, we solve two linear equations. First, an equation on the AF metric (N, g_N), and second, an equation on the compact manifold (Z, g_Z). The "group action" will refer to the $U(2)$-action in the cases g_Z is the Fubini-Study metric and g_N is the Burns metric, and to the toric action plus diagonal symmetry in the case g_Z is $g_{S^2\times S^2}$ and g_N is the corresponding Green's function metric.

7.1. A linear equation on (N, g_N).

On the compact manifold (Z, g_Z), in normal coordinates $\{z^i\}$ around z_0, we have the expansion

$$g_Z = (g_Z)_{ij}dz^i dz^j = (\delta_{ij} + H_2(z)_{ij} + O(|z|^4)_{ij})dz^i dz^j, \tag{7.1}$$

where

$$H_2(z)_{ij} = -\frac{1}{3}R_{ikjl}(z_0)z^k z^l. \tag{7.2}$$

Again let (N, g_N) be the conformal blow-up of the Bach-flat manifold (Y, g_Y), as above. Consider the quadratic tensor

$$H_2(x) = (-\frac{1}{3}R_{ikjl}(z_0)x^k x^l)dx^i dx^j. \tag{7.3}$$

This tensor $H_2(x)$ of course does not live on all of N, since it is only defined in the AF coordinate system. To extend $H_2(x)$ to all of N, let $0 \leq \phi \leq 1$ be a cut-off function satisfying

$$\phi(t) = \begin{cases} 1 & t \leq 1 \\ 0 & t \geq 2, \end{cases} \tag{7.4}$$

and consider $(1 - \phi(R_0^{-1}x))H_2(x)$, where R_0 is very large.

Proposition 7.1. *Let S denote the linearized operator on N, then*

$$S((1 - \phi(R_0^{-1}x)H_2(x)) = O(|x|^{-4}) \tag{7.5}$$

as $|x| \to \infty$

Proof. From (3.43), the linearized operator has the general form

$$\begin{aligned} Sh = (g^{-2} + g * g^{-3}) * \nabla^4 h + g * g^{-3} * Rm * \nabla^2 h + g * g^{-3} * \nabla Rm * \nabla h \\ + (g^{-2} + g * g^{-3}) * (\nabla^2 Rm + Rm * Rm) * h. \end{aligned} \tag{7.6}$$

It is easy to see that for $|x|$ sufficiently large and any tensor h,

$$\begin{aligned} \nabla^4 h = \partial^4 h + \Gamma * \partial^3 h + (\partial\Gamma + \Gamma * \Gamma) * \partial^2 h \\ + (\partial^2\Gamma + \Gamma * \partial\Gamma) * \partial h + (\partial^3\Gamma + \partial\Gamma * \partial\Gamma + \Gamma * \partial^2\Gamma) * h, \end{aligned} \tag{7.7}$$

where ∂ denotes coordinate partial derivatives. If h grows quadratically, then since g_N is AF of order 2, we see that

$$\nabla^4 h = \partial^4 h + O(|x|^{-4}). \tag{7.8}$$

Since $(g_N)_{ij} = \delta_{ij} + O(|x|^{-2})$, it follows that

$$(g^{-2} + g * g^{-3}) * \nabla^4 h = S_0 h + O(|x|^{-4}),$$

where S_0 is the linearized operator with respect to the flat metric. Estimating the other terms on the right-hand side of (7.6) in a similar manner, we find

$$S(h) = S_0 h + O(|x|^{-4}) \tag{7.9}$$

as $|x| \to \infty$. Since H_2 has quadratic leading term and S_0 is a fourth-order operator, we clearly have

$$S_0(H_2) = 0. \tag{7.10}$$

Therefore,

$$S(H_2) = O(|x|^{-4}) \tag{7.11}$$

as $|x| \to \infty$. ☐

Next, given $\epsilon > 0$, consider

$$S : C^{4,\alpha}_{\epsilon}(N) \to C^{0,\alpha}_{\epsilon-4}(N). \tag{7.12}$$

The cokernel of this mapping is the kernel of

$$S^* : C^{4,\alpha}_{-\epsilon}(N) \to C^{0,\alpha}_{-\epsilon-4}(N), \tag{7.13}$$

which consists of the decaying elements.

By Theorem 5.1, $Ker(S^*)$ is 1-dimensional, and spanned by the element

$$o_1 = \mathcal{K}\omega_1 + f \cdot g_N. \tag{7.14}$$

Since $Ker(S^*)$ is nontrivial, this means the map in (7.12) is not surjective, that is, $S(C^{4,\alpha}_{\epsilon}) \subset C^{0,\alpha}_{\epsilon-4}$ is a proper subset, and the quotient space

$$C^{0,\alpha}_{\epsilon-4}/S(C^{4,\alpha}_{\epsilon}) \tag{7.15}$$

is 1-dimensional. A tensor $h \in C^{0,\alpha}_{\epsilon-4}$ is in the image of $C^{4,\alpha}_{\epsilon}$ under S if and only if it pairs trivially with $Ker(S^*)$ under the L^2 pairing. That is

$$h \in S(C^{4,\alpha}_{\epsilon}) \Longleftrightarrow \int_N \langle h, o_1 \rangle dV = 0. \tag{7.16}$$

Since the quotient space is 1-dimensional, we choose $k_1^{(0)} \in C^{0,\alpha}_{\epsilon-4}$ having compact support in $B(x_0, R_0)$ (where x_0 is a basepoint) satisfying

$$\int \langle o_1, k_1^{(0)} \rangle dV = 1, \tag{7.17}$$

and we can write

$$C^{0,\alpha}_{\epsilon-4} = S(C^{4,\alpha}_{\epsilon}) \oplus \mathbb{R} \cdot k_1^{(0)}. \tag{7.18}$$

By averaging over the group, we may assume that $k_1^{(0)}$ is invariant under the group action. Therefore, we can write

$$S((1 - \phi(R_0^{-1}x))H_2(x)) = S(h_\epsilon) + \lambda k_1^{(0)}, \tag{7.19}$$

where $h_\epsilon \in C^{4,\alpha}_{\epsilon}$, and $\lambda \in \mathbb{R}$. Again, by averaging over the group, we may assume that h_ϵ is invariant under the group action. Rewriting this as

$$S((1 - \phi(R_0^{-1}x))H_2 - h_\epsilon) = \lambda k_1^{(0)}, \tag{7.20}$$

we now define

$$\tilde{H}_2 \equiv (1 - \phi(R_0^{-1}x))H_2 - h_\epsilon. \tag{7.21}$$

Since $h_\epsilon \in C^{4,\alpha}_{\epsilon}$, clearly $\tilde{H}_2$ has leading term *exactly* equal to H_2 as $|x| \to \infty$. To summarize, we have solved

Proposition 7.2. *On (N, g_N), there exists a solution of*

$$\begin{aligned} &S(\tilde{H}_2) = \lambda k_1^{(0)}, \\ &\tilde{H}_2(x) = H_2(x) + O^{(4)}(|x|^{\epsilon}), \ \ \textit{as } |x| \to \infty, \end{aligned} \tag{7.22}$$

where $k_1^{(0)}$ *is a tensor with compact support on* (N, g_N) *satisfying*

$$\int_N \langle o_1, k_1^{(0)} \rangle \, dV = 1. \tag{7.23}$$

Furthermore, $\tilde{H}_2$ *can be chosen to be invariant under the group action.*

7.2. **A linear equation on** (Z, g_Z)**.** Next we return to the compact metric (Z, g_Z). Recall on (N, g_N), we have an AF-coordinate system satisfying

$$g_N = (g_N)_{ij} dx^i dx^j = (\delta_{ij} + H_{-2}(x)_{ij} + O(|x|^{-4+\epsilon})_{ij}) dx^i dx^j, \tag{7.24}$$

where

$$H_{-2}(x) = \Big(-\frac{1}{3} R_{ikjl}(y_0) \frac{x^k x^l}{|x|^4} + 2A \frac{1}{|x|^2} \delta_{ij} \Big) dx^i dx^j \tag{7.25}$$

is a 2-tensor with components

$$H_{-2}(x)_{ij} = -\frac{1}{3} R_{ikjl}(y_0) \frac{x^k x^l}{|x|^4} + 2A \frac{1}{|x|^2} \delta_{ij}. \tag{7.26}$$

Consider the inverse quadratic tensor

$$H_{-2}(z) = \Big(-\frac{1}{3} R_{ikjl}(y_0) \frac{z^k z^l}{|z|^4} + 2A \frac{1}{|z|^2} \delta_{ij} \Big) dz^i dz^j. \tag{7.27}$$

Extend this tensor to all of Z by $\phi((R')^{-1} z) H_{-2}(z)$, where $b < R' < inj_{z_0}(g_Z)$ is some fixed radius.

We will need the following technical lemma both in this Section, and later in Section 10:

Lemma 7.3. *Let* S_0 *denote the linearized operator with respect to the flat metric. Then*

$$S_0(H_{-2}) = 0, \tag{7.28}$$

where H_{-2} *is viewed as a tensor on* $\mathbb{R}^4 \setminus \{0\}$*.*

Furthermore, if $(B_0^t)'$ *denotes the linearization of the* B^t*-tensor at the flat metric, then*

$$(B_0^t)'(H_{-2}) = 0. \tag{7.29}$$

Proof. To prove the Lemma we use the expansion (7.24):

$$g_N = g_0 + H_{-2} + O(|x|^{-4+\epsilon}) \tag{7.30}$$

where g_0 is the flat metric. Let $\theta = g_N - g_0$. Since g_N is B^t-flat,

$$\begin{aligned} P_{g_0}(\theta) &= B^t(g_N) + \mathcal{K}_{g_N} \delta_0 \mathcal{K}_0 \delta_0 \overset{\circ}{\theta} \\ &= \mathcal{K}_{g_N} \delta_0 \mathcal{K}_0 \delta_0 \overset{\circ}{\theta}. \end{aligned} \tag{7.31}$$

We can also use the expansion of P at the flat metric g_0 to write

$$\begin{aligned} P_{g_0}(\theta) &= P_{g_0}(0) + S_0(\theta) + Q(\theta) \\ &= B^t(g_0) + S_0(\theta) + Q(\theta) \\ &= S_0(\theta) + Q(\theta). \end{aligned} \tag{7.32}$$

Combining (7.31) and (7.32) we find

$$S_0\theta = \mathcal{K}_{g_N}\delta_0\mathcal{K}_0\delta_0\overset{\circ}{\theta} - Q(\theta). \tag{7.33}$$

Since

$$\theta = H_{-2} + O(|x|^{-4+\epsilon}) \tag{7.34}$$

and S_0 is fourth order,

$$S_0\theta = S_0(H_{-2}) + O(|x|^{-8+\epsilon}), \tag{7.35}$$

hence

$$S_0(H_{-2}) = \mathcal{K}_{g_N}\delta_0\mathcal{K}_0\delta_0\overset{\circ}{\theta} - Q(\theta) + O(|x|^{-8+\epsilon}). \tag{7.36}$$

Also, using (3.27) we have

$$Q(\theta) = O(|x|^{-8}), \tag{7.37}$$

so that

$$S_0(H_{-2}) = \mathcal{K}_{g_N}\delta_0\mathcal{K}_0\delta_0\overset{\circ}{\theta} + O(|x|^{-8+\epsilon}). \tag{7.38}$$

It remains to estimate the gauge-fixing operator acting on θ. By (7.26),

$$\overset{\circ}{\theta}_{ij} = -\frac{1}{3}W_{ikjl}(y_0)\frac{x^k x^l}{|x|^4} + \frac{R(y_0)}{36}\Big\{\frac{x^i x^j}{|x|^4} - \frac{\delta_{ij}}{|x|^2}\Big\} + O(|x|^{-4}). \tag{7.39}$$

Using the skew-symmetry of the Weyl tensor we find

$$(\delta_0\overset{\circ}{\theta})_j = \frac{R(y_0)}{24}\frac{x^j}{|x|^4} + O(|x|^{-5}). \tag{7.40}$$

We next calculate

$$\delta_0\mathcal{K}_0\delta_0\overset{\circ}{\theta} = \Box\delta_0\overset{\circ}{\theta}.$$

It is easy to check that the form

$$\omega = \frac{x^j}{|x|^4}dx^j \tag{7.41}$$

is harmonic. Therefore, using the formula (5.12) for $\Box$ on Euclidean space,

$$\Box\omega = \frac{3}{2}d\delta\omega + \delta d\omega = 0.$$

Consequently,

$$\delta_0\mathcal{K}_0\delta_0\overset{\circ}{\theta} = \Box\delta_0\overset{\circ}{\theta} = \Box\Big(\frac{R(y_0)}{24}\omega + O(|x|^{-5})\Big) = O(|x|^{-7}). \tag{7.42}$$

It follows that

$$\mathcal{K}_{g_N}\delta_0\mathcal{K}_0\delta_0\overset{\circ}{\theta} = O(|x|^{-8}), \tag{7.43}$$

which, using (7.38), implies

$$S_0(H_{-2}) = O(|x|^{-8+\epsilon}). \tag{7.44}$$

However, since H_{-2} is homogeneous of degree -2, $S_0(H_{-2})$ must be homogeneous of degree -6. Therefore, (7.44) implies that $S_0(H_{-2})$ vanishes.

A similar argument (expanding the B^t-tensor as in Proposition 3.8) gives (7.29). $\square$

Proposition 7.4. *Let S denote the linearized operator on Z, then*

$$S(\phi((R')^{-1}z)H_{-2}(z)) = O(|z|^{-4}) \tag{7.45}$$

as $|z| \to 0$.

Proof. As above, for $|z|$ sufficiently small and any tensor h,

$$\begin{aligned}\nabla^4 h &= \partial^4 h + \Gamma * \partial^3 h + (\partial\Gamma + \Gamma * \Gamma) * \partial^2 h \\ &\quad + (\partial^2\Gamma + \Gamma * \partial\Gamma) * \partial h + (\partial^3\Gamma + \partial\Gamma * \partial\Gamma + \Gamma * \partial^2\Gamma) * h,\end{aligned} \tag{7.46}$$

where ∂ denotes coordinates partial derivatives. If h blows-up inverse quadratically, then since $\{z^i\}$ are Riemannian normal coordinates, we see that

$$\nabla^4 h = \partial^4 h + O(|z|^{-4}). \tag{7.47}$$

Arguing as we did in the proof of Proposition 7.1, we find that

$$S(h) = S_0 h + O(|z|^{-4}), \tag{7.48}$$

where S_0 is the linearized operator with respect to the flat metric. If we take $h = H_{-2}$ in (7.48), then (7.28) of Lemma 7.3 gives

$$S(H_{-2}) = O(|z|^{-4}) \tag{7.49}$$

as $|z| \to 0$, and the Proposition follows. $\square$

Next, for $\epsilon > 0$, we have

$$S : C^{4,\alpha}_{-\epsilon}(Z) \to C^{0,\alpha}_{-\epsilon-4}(Z), \tag{7.50}$$

with adjoint mapping

$$S^* : C^{4,\alpha}_{\epsilon}(Z) \to C^{0,\alpha}_{\epsilon-4}(Z). \tag{7.51}$$

By Theorem 4.1, there is no (invariant) cokernel. Thus there exists $h_{-\epsilon} \in C^{4,\alpha}_{-\epsilon}$ such that

$$S(\phi((R')^{-1}z)H_{-2}(z)) = S(h_\epsilon), \tag{7.52}$$

or rather

$$S(\phi((R')^{-1}z)H_{-2}(z) - h_\epsilon) = 0. \tag{7.53}$$

Averaging over the group, we may assume that h_ϵ is invariant under the group action. We then define

$$\tilde{H}_{-2}(z) = \phi((R')^{-1}z)H_{-2}(z) - h_\epsilon. \tag{7.54}$$

To summarize, we have proved

Proposition 7.5. *On (Z, g_Z), there exists a solution $\tilde{H}_{-2}$ of*

$$S(\tilde{H}_{-2}(z)) = 0 \tag{7.55}$$

$$\tilde{H}_{-2}(z) = H_{-2}(z) + O(|z|^{-\epsilon}), \text{ as } |z| \to 0. \tag{7.56}$$

Furthermore, $\tilde{H}_{-2}$ can be chosen to be invariant under the group action.

Remark 7.6. From now on, we will fix $\epsilon > 0$ small.

8. COMPUTATION OF THE LEADING TERM

In this section we compute the constant λ which arose above in Proposition 7.2. As the title of this section indicates, we will refer to this constant as "the leading term" for reasons which will become clear later in Section 12.

Recall from Proposition 7.2 that λ was defined via equation (7.22):

$$S(\tilde{H}_2) = \lambda k_1^{(0)} \text{ on } N, \tag{8.1}$$

with

$$\tilde{H}_2(x) = H_2(x) + O^{(4)}(|x|^\epsilon) \tag{8.2}$$

as $|x| \to \infty$, and

$$(H_2)_{ij} = -\frac{1}{3}R_{ikj\ell}(z_0)x^k x^\ell. \tag{8.3}$$

Pairing both side of the defining equation for λ with the cokernel element o_1 and integrating gives

$$\lambda = \int_N S(\tilde{H}_2), o_1\rangle \, dV, \tag{8.4}$$

since

$$\int \langle k_1^{(0)}, o_1\rangle \, dV = 1.$$

Proposition 8.1. *The constant λ is given by*

$$\lambda = \frac{4}{9}\omega_3\big[W_{ikj\ell}(y_0)W_{ikj\ell}(z_0) + W_{ikj\ell}(y_0)W_{i\ell jk}(z_0)\big] + 4t\omega_3 R(z_0)\mathrm{mass}(g_N), \tag{8.5}$$

where $\omega_3 = Vol(S^3)$.

We prove this formula through a series of lemmas. To begin, let

$$B = \{x \in N \ : \ |x| < a^{-1}\}, \tag{8.6}$$

(where we extend $|x|$ to be defined on all of N by letting it be a constant outside of the AF region of N), and use (3.15) to write

$$\begin{aligned}\int_B \langle S\tilde{H}_2, o_1\rangle &= \\ &\int_B \langle \Delta^2(\overset{\circ}{\tilde{H}}_2), o_1\rangle + \int_B \langle \mathcal{K}[d(\mathcal{D}_2(\tilde{H}_2))], o_1\rangle + \frac{3}{2}t\int_B \langle \big[\Delta^2(tr\ \tilde{H}_2) - \Delta(\delta^2 \tilde{H}_2)\big] g, o_1\rangle,\end{aligned} \tag{8.7}$$

where $\overset{\circ}{T}$ denotes the trace-free part of the symmetric two-tensor T.

Lemma 8.2. *As $a \to 0$,*

$$\begin{aligned}\int_B \langle \Delta^2 \overset{\circ}{\tilde{H}}_2, o_1\rangle = \int_B &\langle \tilde{H}_2, \Delta^2 \kappa\rangle \\ &+ \frac{4}{9}\omega_3\big[W_{ikj\ell}(y_0)W_{ikj\ell}(z_0) + W_{ikj\ell}(y_0)W_{i\ell jk}(z_0)\big] + o(1).\end{aligned} \tag{8.8}$$

Proof. Since κ is the trace-free part of o_1,

$$\int_B \langle \Delta^2 \overset{\circ}{\tilde{H}}_2, o_1\rangle = \int_B \langle \Delta^2 \overset{\circ}{\tilde{H}}_2, \kappa\rangle.$$

Integrating by parts,

$$\begin{aligned}\int_B \langle \Delta^2 \overset{\circ}{\tilde{H}}_2, \kappa\rangle = \int_B \langle \overset{\circ}{\tilde{H}}_2, \Delta^2\kappa\rangle &+ \oint_{\partial B} \langle \nabla_N(\Delta \overset{\circ}{\tilde{H}}_2), \kappa\rangle - \oint_{\partial B} \langle \Delta \overset{\circ}{\tilde{H}}_2, \nabla_N \kappa\rangle \\ &+ \oint_{\partial B} \langle \nabla_N \overset{\circ}{\tilde{H}}_2, \Delta\kappa\rangle - \oint_{\partial B} \langle \overset{\circ}{\tilde{H}}_2, \nabla_N(\Delta\kappa)\rangle,\end{aligned} \tag{8.9}$$

where N is the outward unit normal to N. All the boundary integrals in (8.9) are with respect to the approximate metric g. To estimate each boundary term we use the fact that on ∂B, the metric and Christoffel symbols satisfy

$$\begin{aligned} g &= \delta + O(a^2), \\ \Gamma &= O(a^3), \\ \partial\Gamma &= O(a^4), \end{aligned} \tag{8.10}$$

where δ denotes the flat metric. For a symmetric 2-tensor $T = T_{ij}$,

$$\Delta T = g^{\alpha\beta}\nabla_\alpha \nabla_\beta T, \tag{8.11}$$

and

$$\nabla_\alpha \nabla_\beta T = \partial_\alpha \partial_\beta T + \Gamma * \partial T + \partial\Gamma * T + \Gamma * \Gamma * T, \tag{8.12}$$

hence

$$\Delta T_{ij} = \Delta_0 T_{ij} + O(a^2) * \partial^2 T + O(a^3) * \partial T + O(a^4) * T, \tag{8.13}$$

where Δ_0 denotes the flat Laplacian.

Taking $T = \overset{\circ}{\tilde{H}}_2$ and using (8.2) we first note

$$\begin{aligned}(\overset{\circ}{\tilde{H}}_2)_{ij} &= (\tilde{H}_2)_{ij} - \frac{1}{4}\big[g^{\alpha\beta}(\tilde{H}_2)_{\alpha\beta}\big]g_{ij}\\ &= -\frac{1}{3}R_{ikj\ell}(z_0)x^k x^\ell + \frac{1}{12}R_{k\ell}(z_0)x^k x^\ell \delta_{ij} + O(|x|^\epsilon).\end{aligned} \tag{8.14}$$

Therefore,

$$\begin{aligned}\partial_\alpha(\overset{\circ}{\tilde{H}}_2)_{ij} &= -\frac{1}{3}R_{i\alpha j\ell}(z_0)x^\ell - \frac{1}{3}R_{ikj\alpha}(z_0)x^k + \frac{1}{6}R_{\alpha k}(z_0)x^k\delta_{ij} + O(|x|^{\epsilon-1}),\\ \partial_\alpha\partial_\beta(\overset{\circ}{\tilde{H}}_2)_{ij} &= -\frac{1}{3}R_{i\alpha j\beta}(z_0) - \frac{1}{3}R_{i\beta j\alpha}(z_0) + \frac{1}{6}R_{\alpha\beta}(z_0)\delta_{ij} + O(|x|^{\epsilon-2}),\end{aligned} \tag{8.15}$$

hence

$$(\Delta\overset{\circ}{\tilde{H}}_2)_{ij} = -\frac{2}{3}\big[R_{ij}(z_0) - \frac{1}{4}R(z_0)\delta_{ij}\big] + O(|x|^{\epsilon-2}). \tag{8.16}$$

Assuming (Z, g_Z) is Einstein, then

$$R_{ij}(z_0) - \frac{1}{4}R(z_0)\delta_{ij} = 0.$$

It follows that

$$\begin{aligned}(\Delta\overset{\circ}{\tilde{H}}_2)_{ij} &= O(|x|^{\epsilon-2}),\\ \nabla_N(\Delta\overset{\circ}{\tilde{H}}_2)_{ij} &= O(|x|^{\epsilon-3}),\end{aligned} \tag{8.17}$$

as $|x| \to \infty$.

By (6.4), on ∂B we have

$$\begin{aligned}|\kappa| &= O(a^2),\\ |\nabla\kappa| &= O(a^3).\end{aligned} \tag{8.18}$$

Using these estimates along with those of (8.17) we find

$$\begin{aligned}\Big|\oint_{\partial B}\langle\nabla_N(\Delta\overset{\circ}{\tilde{H}}_2), \kappa\rangle\Big| &= O(a^{2-\epsilon}),\\ \Big|\oint_{\partial B}\langle\Delta\overset{\circ}{\tilde{H}}_2, \nabla_N\kappa\rangle\Big| &= O(a^{2-\epsilon}).\end{aligned} \tag{8.19}$$

Next, we take $T = \kappa_{ij}$. Using (6.4),

$$\begin{aligned}
\partial_\alpha \kappa_{ij} &= \frac{2}{3} W_{i\alpha j\ell}(y_0) \frac{x^\ell}{|x|^4} + \frac{2}{3} W_{ikj\alpha}(y_0) \frac{x^k}{|x|^4} - \frac{8}{3} W_{ikj\ell}(y_0) \frac{x^k x^\ell x^\alpha}{|x|^6} + O(a^4), \\
\partial_\alpha \partial_\beta \kappa_{ij} &= \frac{2}{3} W_{i\alpha j\beta}(y_0) \frac{1}{|x|^4} + \frac{2}{3} W_{i\beta j\alpha}(y_0) \frac{1}{|x|^4} + 16 W_{ikj\ell}(y_0) \frac{x^k x^\ell x^\alpha x^\beta}{|x|^8} \\
&\quad - \frac{8}{3} W_{i\alpha j\ell}(y_0) \frac{x^\ell x^\beta}{|x|^6} - \frac{8}{3} W_{ikj\alpha}(y_0) \frac{x^k x^\beta}{|x|^6} - \frac{8}{3} W_{i\beta j\ell}(y_0) \frac{x^\ell x^\alpha}{|x|^6} \\
&\quad - \frac{8}{3} W_{ikj\beta}(y_0) \frac{x^k x^\alpha}{|x|^6} - \frac{8}{3} W_{ikj\ell}(y_0) \frac{x^k x^\ell}{|x|^6} \delta_{\alpha\beta} + O(a^5).
\end{aligned} \tag{8.20}$$

Therefore,

$$\begin{aligned}
\Delta \kappa_{ij} &= -\frac{16}{3} W_{ikj\ell}(y_0) \frac{x^k x^\ell}{|x|^6} + O(a^5) \\
&= -\frac{8}{|x|^2} \kappa_{ij} + O(a^5).
\end{aligned} \tag{8.21}$$

On ∂B,

$$N^\alpha = \frac{x^\alpha}{|x|} + O(a^2), \tag{8.22}$$

hence

$$\nabla_N T_{ij} = \frac{x^\alpha}{|x|} \partial_\alpha T_{ij} + O(a^2) * \partial T + O(a^3) * T. \tag{8.23}$$

From (8.14),(8.20), and (8.23) (or, by reasons of homogeneity) we conclude

$$\begin{aligned}
\nabla_N (\mathring{\tilde{H}}_2)_{ij} &= -\frac{2}{3} R_{ikj\ell}(z_0) \frac{x^k x^\ell}{|x|} + \frac{1}{6} R_{k\ell}(z_0) \frac{x^k x^\ell}{|x|} \delta_{ij} + O(a^{1-\epsilon}) \\
&= \frac{2}{|x|} (\mathring{\tilde{H}}_2)_{ij} + O(a^{1-\epsilon}), \\
\nabla_N (\Delta \kappa)_{ij} &= \frac{64}{3} W_{ikj\ell}(y_0) \frac{x^k x^\ell}{|x|^7} + O(a^7) \\
&= \frac{32}{|x|^3} \kappa_{ij} + O(a^7).
\end{aligned} \tag{8.24}$$

It follows that

$$\langle \nabla_N \mathring{\tilde{H}}_2, \Delta \kappa \rangle - \langle \mathring{\tilde{H}}_2, \nabla_N (\Delta \kappa) \rangle = -48 \frac{1}{|x|^3} \langle \mathring{\tilde{H}}_2, \kappa \rangle + O(a^4), \tag{8.25}$$

hence

$$\oint_{\partial B} \langle \nabla_N \mathring{\tilde{H}}_2, \Delta \kappa \rangle - \langle \mathring{\tilde{H}}_2, \nabla_N (\Delta \kappa) \rangle = \frac{32}{3} \int_{|\xi|=1} W_{ikj\ell}(y_0) R_{i\alpha j\beta}(z_0) \xi^k \xi^\ell \xi^\alpha \xi^\beta \, dS + O(a). \tag{8.26}$$

If we decompose the curvature tensor of $R_{i\alpha j\beta}(z_0)$ (again assuming (Z, g_Z) is Einstein),

$$R_{i\alpha j\beta}(z_0) = W_{i\alpha j\beta}(z_0) + \frac{1}{12}R(z_0)\big(\delta_{ij}\delta_{\alpha\beta} - \delta_{i\beta}\delta_{j\alpha}\big).$$

Therefore, the integrand in (8.26) can be written

$$W_{ikj\ell}(y_0)R_{i\alpha j\beta}(z_0) = W_{ikj\ell}(y_0)W_{i\alpha j\beta}(z_0) - \frac{1}{12}R(z_0)W_{\beta k\alpha\ell}(y_0).$$

hence

$$\begin{aligned}\int_{|\xi|=1} W_{ikj\ell}(y_0)R_{i\alpha j\beta}(z_0)\xi^k\xi^\ell\xi^\alpha\xi^\beta\ dS = \int_{|\xi|=1} & W_{ikj\ell}(y_0)W_{i\alpha j\beta}(z_0)\xi^k\xi^\ell\xi^\alpha\xi^\beta\ dS \\ & - \frac{1}{12}R(z_0)\int_{|\xi|=1} W_{\beta k\alpha\ell}(y_0)\xi^k\xi^\ell\xi^\alpha\xi^\beta\ dS.\end{aligned} \tag{8.27}$$

The last integral vanishes by skew-symmetry of the Weyl tensor; therefore,

$$\oint_{\partial B}\langle\nabla_N \mathring{\tilde{H}}_2, \Delta\kappa\rangle - \langle\mathring{\tilde{H}}_2, \nabla_N(\Delta\kappa)\rangle = \frac{32}{3}\int_{|\xi|=1} W_{ikj\ell}(y_0)W_{i\alpha j\beta}(z_0)\xi^k\xi^\ell\xi^\alpha\xi^\beta\ dS + O(a). \tag{8.28}$$

We now use the identity (see [Bre08])

$$\int_{|\xi|=1}\xi^k\xi^\ell\xi^\alpha\xi^\beta\ dS = \frac{\omega_3}{24}\big(\delta_{k\ell}\delta_{\alpha\beta} + \delta_{k\alpha}\delta_{\beta\ell} + \delta_{k\beta}\delta_{\alpha\ell}\big). \tag{8.29}$$

Plugging this into (8.28), we obtain

$$\begin{aligned}\oint_{\partial B}\langle\nabla_N \mathring{\tilde{H}}_2, \Delta\kappa\rangle - \langle\mathring{\tilde{H}}_2, \nabla_N(\Delta\kappa)\rangle = \frac{4}{9}\omega_3\big[& W_{ikj\ell}(y_0)W_{ikj\ell}(z_0) + W_{ikj\ell}(y_0)W_{i\ell jk}(z_0)\big] \\ & + O(a),\end{aligned} \tag{8.30}$$

which proves the Lemma. □

Lemma 8.3. *As $a \to 0$,*

$$\int_B\langle\mathcal{K}[d(\mathcal{D}_2(\tilde{H}_2)], o_1\rangle = O(a^{2-\epsilon}). \tag{8.31}$$

Proof. Since $\mathcal{K}[\cdot]$ is trace-free, we can rewrite the integrand in (8.31) as

$$\int_B\langle\mathcal{K}[d(\mathcal{D}_2(\tilde{H}_2)], o_1\rangle = \int_B\langle\mathcal{K}[d(\mathcal{D}_2(\tilde{H}_2)], \kappa\rangle.$$

Integrating by parts and using the fact that κ is divergence-free, we get

$$\begin{aligned}\int_B \langle \mathcal{K}[d(\mathcal{D}_2(\tilde{H}_2)], \kappa\rangle &= -2\int_B \langle d(\mathcal{D}_2(\tilde{H}_2), \delta\kappa\rangle \\ &\quad + 2\oint_{\partial B} \kappa(d(\mathcal{D}_2(\tilde{H}_2)), N) \\ &= 2\oint_{\partial B} \kappa(d(\mathcal{D}_2(\tilde{H}_2)), N).\end{aligned} \tag{8.32}$$

Using (8.12) and computing as we did in the proof of Lemma 8.2, on ∂B we find

$$\begin{aligned}\delta^2\tilde{H}_2 &= \frac{1}{3}R(z_0) + O(a^{2-\epsilon}), \\ \Delta(tr\ \tilde{H}_2) &= -\frac{2}{3}R(z_0) + O(a^{2-\epsilon}).\end{aligned} \tag{8.33}$$

Therefore,

$$\begin{aligned}\mathcal{D}_2(\tilde{H}_2) &= (t + \frac{5}{12})R(z_0) + O(a^{2-\epsilon}), \\ d(\mathcal{D}_2(\tilde{H}_2)) &= O(a^{3-\epsilon}).\end{aligned}$$

Since $\kappa = O(a^2)$ on ∂B, we see that the boundary term in (8.32) is $O(a^{2-\epsilon})$, which proves the Lemma. ☐

Lemma 8.4. *As $a \to 0$,*

$$\begin{aligned}\frac{3}{2}t\int_B \langle \big[\Delta^2(tr\ \tilde{H}_2) - \Delta(\delta^2\tilde{H}_2)\big]g, o_1\rangle &= \int_B \langle \tilde{H}_2, 6t\big[(\Delta^2 f)g - \nabla^2(\Delta f)\big]\rangle \\ &\quad + 4t\Big(12A - \frac{R(y_0)}{12}\Big)\omega_3 R(z_0) + o(1).\end{aligned} \tag{8.34}$$

Proof. Since $o_1 = \kappa + fg$ with κ trace-free, we have

$$\begin{aligned}\frac{3}{2}t\int_B \langle \big[\Delta^2(tr\ \tilde{H}_2) - \Delta(\delta^2\tilde{H}_2)\big]g, o_1\rangle &= \frac{3}{2}t\int_B \langle \big[\Delta^2(tr\ \tilde{H}_2) - \Delta(\delta^2\tilde{H}_2)\big]g, \kappa + fg\rangle \\ &= \frac{3}{2}t\int_B \langle \big[\Delta^2(tr\ \tilde{H}_2) - \Delta(\delta^2\tilde{H}_2)\big]g, fg\rangle \\ &= 6t\int_B \big[\Delta^2(tr\ \tilde{H}_2) - \Delta(\delta^2\tilde{H}_2)\big]f.\end{aligned} \tag{8.35}$$

Integrating by parts, we find

$$\begin{aligned}
\int_B \big[\Delta^2(tr\ \tilde{H}_2) - \Delta(\delta^2\tilde{H}_2)\big] f &= \int_B \langle \tilde{H}_2, (\Delta^2 f)g - \nabla^2(\Delta f)\rangle \\
&+ \oint_{\partial B} f\frac{\partial}{\partial N}\Delta(tr\ \tilde{H}_2) - \oint_{\partial B} \Delta(tr\ \tilde{H}_2)\frac{\partial}{\partial N} f + \oint_{\partial B} \frac{\partial}{\partial N}(tr\ \tilde{H}_2)\Delta f \\
&- \oint_{\partial B} (tr\ \tilde{H}_2)\frac{\partial}{\partial N}(\Delta f) + \oint_{\partial B} \tilde{H}_2(N, \nabla(\Delta f)) - \oint_{\partial B} (\Delta f)\langle \delta\tilde{H}_2, N\rangle \\
&+ \oint_{\partial B} \frac{\partial}{\partial N} f(\delta^2\tilde{H}_2) - \oint_{\partial B} f\frac{\partial}{\partial N}(\delta^2\tilde{H}_2) \\
&= \int_B \langle \tilde{H}_2, (\Delta^2 f)g - \nabla^2(\Delta f)\rangle + I_1 + \cdots + I_8.
\end{aligned} \tag{8.36}$$

By Theorem 5.1, on ∂B

$$\begin{aligned}
\Delta f &= -\frac{1}{3}\langle Ric, o_1\rangle = O(a^6), \\
\frac{\partial}{\partial N}(\Delta f) &= -\frac{1}{3}\langle \nabla_N Ric, o_1\rangle - \frac{1}{3}\langle Ric, \nabla_N o_1\rangle = O(a^7).
\end{aligned} \tag{8.37}$$

Also, from the preceding lemma (see (8.33))

$$\begin{aligned}
tr\ \tilde{H}_2 &= -\frac{1}{3}R_{k\ell}(z_0)x^k x^\ell + O(a^{-\epsilon}) = O(a^{-2}), \\
\frac{\partial}{\partial N}(tr\ \tilde{H}_2) &= -\frac{2}{3}R_{k\ell}(z_0)\frac{x^k x^\ell}{|x|} + O(a^{1-\epsilon}) = O(a^{-1}), \\
\delta^2\tilde{H}_2 &= \frac{1}{3}R(z_0) + O(a^{2-\epsilon}) = O(1), \\
\Delta(tr\ \tilde{H}_2) &= -\frac{2}{3}R(z_0) + O(a^{2-\epsilon}) = O(1) \\
\frac{\partial}{\partial N}(\delta^2\tilde{H}_2) &= O(a^{3-\epsilon}), \\
\frac{\partial}{\partial N}(\Delta(tr\ \tilde{H}_2)) &= O(a^{3-\epsilon}).
\end{aligned} \tag{8.38}$$

Therefore,

$$
\begin{aligned}
I_1 &= \oint_{\partial B} f \frac{\partial}{\partial N} \Delta (tr\ \tilde{H}_2) = O(a^{2-\epsilon}) \\
I_3 &= \oint_{\partial B} \frac{\partial}{\partial N} (tr\ \tilde{H}_2) \Delta f = O(a^2), \\
I_4 &= -\oint_{\partial B} (tr\ \tilde{H}_2) \frac{\partial}{\partial N} (\Delta f) = O(a^2), \\
I_5 &= \oint_{\partial B} \tilde{H}_2(N, \nabla(\Delta f)) = O(a^2), \\
I_6 &= -\oint_{\partial B} (\Delta f) \langle \delta \tilde{H}_2, N \rangle = O(a^2), \\
I_8 &= -\oint_{\partial B} f \frac{\partial}{\partial N} (\delta^2 \tilde{H}_2) = O(a^{2-\epsilon}).
\end{aligned}
\tag{8.39}
$$

Therefore, it remains to calculate I_2 and I_7.

First, using (8.38) we have

$$
\begin{aligned}
I_2 + I_7 &= \oint_{\partial B} \left[\delta^2 \tilde{H}_2 - \Delta (tr\ \tilde{H}_2) \right] \frac{\partial}{\partial N} f \\
&= \oint_{\partial B} \left[R(z_0) + O(a^{2-\epsilon}) \right] \frac{\partial}{\partial N} f \\
&= R(z_0) \oint_{\partial B} \frac{\partial}{\partial N} f + O(a^{2-\epsilon}) \quad \text{(by (5.6))} \\
&= R(z_0) \int_B \Delta f + O(a^{2-\epsilon}) \ \ \text{(by the divergence theorem)} \\
&= -\frac{1}{3} R(z_0) \int_B \langle Ric, \kappa \rangle + O(a^{2-\epsilon}) \ \ \text{(by (5.5))}.
\end{aligned}
\tag{8.40}
$$

Using the fact that $\kappa = \mathcal{K}[\omega_1]$, we can integrate by parts to obtain

$$
\begin{aligned}
\int_B \langle Ric, \kappa \rangle &= \int_B \langle Ric, \mathcal{K}[\omega_1] \rangle \\
&= 2 \int_B R_{ij} \nabla^i \omega_1^j \\
&= -2 \int_B \nabla^i R_{ij} \omega_1^j + 2 \oint_{\partial B} R_{ij} N^i \omega_1^j
\end{aligned}
$$

Using the second Bianchi identity and the fact that the scalar curvature is zero, the solid integral above vanishes and we conclude

$$
I_2 + I_7 = -\frac{2}{3} R(z_0) \oint_{\partial B} Ric(N, \omega_1) + O(a^{2-\epsilon}). \tag{8.41}
$$

By (5.6) and (8.22),

$$Ric(N, \omega_1) = R_{ij}\frac{x^i x^j}{|x|} + O(|x|^{-5}). \tag{8.42}$$

Proposition 8.5. *As* $|x| \to \infty$*,*

$$\begin{aligned} R_{ij} = &-\frac{4}{3}W_{ikj\ell}(y_0)\frac{x^k x^\ell}{|x|^6} - \frac{1}{36}R(y_0)\frac{1}{|x|^4}\delta_{ij} + \frac{1}{9}R_(y_0)\frac{x^i x^j}{|x|^6} \\ &- \frac{16A}{|x|^6}x^i x^j + \frac{4A}{|x|^4}\delta_{ij} + O(|x|^{-5}). \end{aligned} \tag{8.43}$$

Proof. This is proved in Appendix A. □

Assuming the proposition, we see that

$$R_{ij}\frac{x^i x^j}{|x|} = \frac{1}{12}R(y_0)\frac{1}{|x|^3} - \frac{12A}{|x|^3} + O(|x|^{-4}).$$

Therefore,

$$\begin{aligned} I_2 + I_7 &= -\frac{2}{3}R(z_0)\oint_{\partial B} Ric(N, \omega_1) + O(a^{2-\epsilon}) \\ &= -\frac{2}{3}R(z_0)\big[\frac{1}{12}R(y_0) - 12A\big]\oint_{\partial B}\frac{1}{|x|^3} + O(a) \\ &= -\frac{2}{3}\omega_3 R(z_0)\big[\frac{1}{12}R(y_0) - 12A\big] + O(a). \end{aligned}$$

Plugging this into (8.35) and (8.36), we arrive at (8.34). □

Combining Lemmas 8.2, 8.3, and 8.4, and using (8.7), we have

$$\begin{aligned} \int_B \langle S\tilde{H}_2, o_1\rangle = &\int_B \langle \tilde{H}_2, \Delta^2\kappa + 6t\big[(\Delta^2 f)g - \nabla^2(\Delta f)\big]\rangle \\ &+ \frac{4}{9}\omega_3\big[W_{ikj\ell}(y_0)W_{ikj\ell}(z_0) + W_{ikj\ell}(y_0)W_{i\ell jk}(z_0)\big] \\ &+ 4t\omega_3 R(z_0)\big\{12A - \frac{1}{12}R(y_0)\big\} + o(1). \end{aligned} \tag{8.44}$$

By Proposition 2.5, the quantity is braces is exactly the mass of the AF space. Proposition 8.1 then follows from the next Lemma:

Lemma 8.6. *The cokernel element* o_1 *satisfies*

$$0 = S(o_1) = \Delta^2\kappa + 6t\big[(\Delta^2 f)g - \nabla^2(\Delta f)\big] + O(|x|^{-8}). \tag{8.45}$$

Proof. By the formula in (3.15), we have

$$So_1 = \Delta^2[o_1 - \frac{1}{4}(tr\ o_1)g] + \mathcal{K}\big[d(\mathcal{D}_2(o_1))\big] + \frac{3}{2}t\Big[\Delta^2(tr\ o_1) - \Delta(\delta^2 o_1)\Big]g + \cdots \tag{8.46}$$

Using the properties of o_1 in Theorem 5.1, we have

$$\begin{aligned} o_1 - \frac{1}{4}(tr\ o_1)g &= \kappa, \\ tr\ o_1 &= 4f, \\ \delta^2 o_1 &= \Delta f, \\ \Delta(tr\ o_1) &= 4\Delta f. \end{aligned}$$

It follows that

$$\begin{aligned} \mathcal{D}_2(o_1) &= \big(t + \frac{5}{6}\big)\delta^2 o_1 - \big(t + \frac{5}{24}\big)\Delta(tr\ o_1) \\ &= -3t\Delta f, \end{aligned}$$

and

$$\begin{aligned} \mathcal{K}\big[d(\mathcal{D}_2(o_1))\big] &= -3t\mathcal{K}\big[d(\Delta f)\big] \\ &= -6t\nabla^2(\Delta f) + \frac{3}{2}t(\Delta^2 f)g \end{aligned}$$

Therefore,

$$\begin{aligned} So_1 &= \Delta^2\kappa + \mathcal{K}\big[d(\mathcal{D}_2(o_1))\big] + \frac{9}{2}t(\Delta^2 f)g + \cdots \\ &= \Delta^2\kappa - 6t\nabla^2(\Delta f) + \frac{3}{2}t(\Delta^2 f)g + \frac{9}{2}t(\Delta^2 f)g + \cdots \\ &= \Delta^2\kappa + 6t\big[(\Delta^2 f)g - \nabla^2(\Delta f)\big] + \cdots, \end{aligned}$$

as claimed. □

9. Naïve approximate metric

Let (Z, g_Z) be a compact B^t-flat manifold. In our application, (Z, g_Z) will be taken to be either $\mathbb{CP}^2$ with the Fubini-Study metric, or $S^2 \times S^2$ with the product metric, with the coordinate systems described in Subsections 2.1 and 2.3.

We let z_0 denote the base point, which is $[1, 0, 0]$ in the case of $\mathbb{CP}^2$, or (n, n) in the case of $S^2 \times S^2$. As seen above, we have a Riemannian normal coordinate system $\{z^i\}, i = 1 \dots 4$, satisfying

$$g_Z = dz^2 + \eta_Z(z), \tag{9.1}$$

where η_Z has the expansion $\eta_Z = (\eta_Z(z))_{ij} dz^i dz^j$ with

$$(\eta_Z(z))_{ij} = -\frac{1}{3}R_{ikjl}(z_0)z^k z^l + O(|z|^4) \tag{9.2}$$

as $|z| \to 0$.

Furthermore, in the case of $\mathbb{CP}^2$ the metric is invariant under the standard linear action of $U(2)$ in the $\{z\}$-coordinates, and in the case of $S^2 \times S^2$ the metric is invariant under the standard diagonal torus action, and also invariant under the diagonal symmetry, both in the $\{z\}$-coordinates.

Next, let (N, g_N) be a B^t-flat AF space of order 2. In our application (N, g_N) will be taken to the either the Burns metric or Green's function metric of the product metric with AF coordinate system as described in Subsections 2.2 and 2.4. The Green's function here is with respect to the basepoint which we will denote as y_0, which is $[1, 0, 0]$ in the case of $\mathbb{CP}^2$, or either point (n, n) of $S^2 \times S^2$.

We denote the AF coordinates as $\{x^i\}, i = 1 \dots 4$, and write

$$g_N = dx^2 + \eta_N, \tag{9.3}$$

where the tensor η_N admits the expansion

$$(\eta_N)_{ij}(x) = -\frac{1}{3} R_{ikjl}(y_0) \frac{x^k x^l}{|x|^4} + 2A \frac{1}{|x|^2} \delta_{ij} + O(|x|^{-4+\epsilon}) \tag{9.4}$$

as $|x| \to \infty$.

In the case of the Burns metric, the metric is invariant under the standard linear action of $U(2)$ in the $\{x\}$-coordinates, and in the case of the Green's function metric on $S^2 \times S^2$, the metric is invariant under the standard diagonal torus action, and also invariant under the diagonal symmetry, both in the $\{x\}$-coordinates.

Let ϕ be the cutoff function defined in (7.4):

$$\phi(t) = \begin{cases} 1 & t \leq 1 \\ 0 & t \geq 2. \end{cases} \tag{9.5}$$

For $b > 0$ denote the annulus $A_Z(b, 2b) = \{b \leq |z| \leq 2b\} \subset Z$, and for $a > 0$ denote the annulus $A_N(a^{-1}, 2a^{-1}) \equiv \{a^{-1} \leq |x| \leq 2a^{-1}\} \subset N$. Let $\iota : A_N(a^{-1}, 2a^{-1}) \to A_Z(b, 2b)$ denote the map $\iota(x) = abx = z$. Identify the annular region $A_Z(b, 2b) \subset X$ with $A_N(a^{-1}, 2a^{-1}) \subset N$ using the map ι to define a new manifold $X_{a,b}$.

Remark 9.1. With this choice of ι, the manifold $X_{a,b}$ is diffeomorphic to $X \# \overline{N_c}$, where N_c is the one-point compactification of N. If we instead choose ι to be defined by, for example, $\iota(x_1, x_2, x_3, x_4) = ab(-x_1, x_2, x_3, x_4)$, $X_{a,b}$ will be diffeomorphic to $X \# N_c$, which can be different topologically.

In the case where (Z, g_Z) is the Fubini-Study metric and (N, g_N) is the Burns metric, the $U(2)$ action extends to $X_{a,b}$, since the actions agree in the coordinate systems. In all other cases, the torus action as well as the diagonal symmetry extend to actions on $X_{a,b}$. For convenience, we will now refer to this action as "the group action", keeping in mind that the group depends on the example.

We compute that

$$\begin{aligned} \iota^*(a^{-2} b^{-2} g_Z) &= a^{-2} b^{-2} \{a^2 b^2 dx^2 + (\iota^* \eta_Z)(x)\} \\ &= dx^2 + \tilde{\eta}_Z(x), \end{aligned} \tag{9.6}$$

where

$$\tilde{\eta}_Z(x) = a^{-2} b^{-2} (\iota^* \eta_Z)(x). \tag{9.7}$$

Note that $\tilde{\eta}_Z$ admits the expansion $\tilde{\eta}_Z = (\tilde{\eta}_Z(x))_{ij} dx^i dx^j$ with

$$(\tilde{\eta}_Z(x))_{ij} = -\frac{1}{3} a^2 b^2 R_{ikjl}(z_0) x^k x^l + \text{ (higher)} \tag{9.8}$$

as $b \to 0$ and for $x \in A_N(a^{-1}, 2a^{-1})$.

Define a metric $g^{(0)}_{a,b}$ on $X_{a,b}$ by

$$g^{(0)}_{a,b} = \begin{cases} a^{-2}b^{-2} g_Z & |z| \geq 2b \\ dx^2 + \phi(a|x|)\eta_N(x) + [1 - \phi(a|x|)]\tilde{\eta}_Z(x) & a^{-1} \leq |x| \leq 2a^{-1} \\ g_N & |x| \leq a^{-1}. \end{cases} \tag{9.9}$$

The group action is linear in the $\{x\}$-coordinates, and is contained in $SO(4)$. Since the cutoff function is radial, it is clear that $g^{(0)}_{a.b}$ is invariant under the group action.

On the damage zone $A_N(a^{-1}, 2a^{-1})$, we will also write the metric as

$$g^{(0)}_{a,b} = dx^2 + \eta_1 + \eta_2, \tag{9.10}$$

where

$$\begin{aligned} \eta_1(x) &= \phi(a|x|)\eta_N(x) \\ \eta_2(x) &= [1 - \phi(a|x|)]\tilde{\eta}_Z(x). \end{aligned} \tag{9.11}$$

Notice that after scaling and identifying, we have

$$|Rm(\iota^*(a^{-2}b^{-2}g_Z^b))| = \begin{cases} a^2b^2(|R(g_Z)| \circ \iota) & |x| \geq 2a^{-1} \\ O(a^2b^2) & a^{-1} \leq |x| \leq 2a^{-1} \\ 0 & |x| \leq 2a^{-1}. \end{cases} \tag{9.12}$$

This implies that

$$|B^t(\iota^*(a^{-2}b^{-2}g_Z^b))| = \begin{cases} 0 & |x| \geq 2a^{-1} \\ O(a^4b^2) & a^{-1} \leq |x| \leq 2a^{-1} \\ 0 & |x| \leq 2a^{-1}. \end{cases} \tag{9.13}$$

Consequently,

$$|B^t(g^{(0)}_{a,b})| = \begin{cases} 0 & |x| \geq 2a^{-1} \\ O(a^4b^2) + O(a^6) & a^{-1} \leq |x| \leq 2a^{-1} \\ 0 & |x| \leq 2a^{-1}, \end{cases} \tag{9.14}$$

which is proved by using the expansion

$$B^t(g^{(0)}_{a,b}) = B^t(g_0) + (B^t)'_{g_0}(\eta_1 + \eta_2) + \mathcal{Q}(\eta_1 + \eta_2), \tag{9.15}$$

where $g_0 = dx^2$ in the damage zone; see Remark 3.8.

This estimate will not suffice for our purposes, and in Section 10 we will construct a “better” approximate metric.

9.1. **Gluing with one basepoint.** To summarize, $(X_{a,b}, g^{(0)}_{a,b})$ is defined in the following cases:

- (i) $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$; the Fubini-Study metric with a Burns metric attached at one fixed point. This case admits a $U(2)$-action.
- (ii) $S^2 \times S^2 \# \overline{\mathbb{CP}}^2 = \mathbb{CP}^2 \# 2\overline{\mathbb{CP}}^2$; the product metric on $S^2 \times S^2$ with a Burns metric attached at one fixed point. Alternatively, we can view this as the Fubini-Study metric on $\mathbb{CP}^2$, with a Green's function $S^2 \times S^2$ metric attached at one fixed point. For this topology, we will therefore construct two different critical metrics.
- (iii) $2\# S^2 \times S^2$; the product metric on $S^2 \times S^2$ with a Green's function $S^2 \times S^2$ metric attached at one fixed point.

All of these cases are invariant under the torus action, and invariant under the diagonal symmetry.

As mentioned in the introduction, the product metric on $S^2 \times S^2$ admits the Einstein quotient $S^2 \times S^2/\mathbb{Z}_2$, where $\mathbb{Z}_2$ acts by the antipodal map on both factors, and the quotient $\mathbb{RP}^2 \times \mathbb{RP}^2$. The diagonal symmetry clearly extends to these metrics. Using one of these metrics as the compact factor or the Green's function metric of one of these as the AF space, we obtain approximate metrics on the non-simply-connected topologies listed in Table 1.3. Note that in this table, the first special value of t_0 corresponds to the the first factor being the compact factor, and the second factor being the AF space, while the second value of t_0 corresponds to the reverse. From this, the approximate metric is clear and we need not detail every case here.

9.2. **Gluing with multiple bubbles.** We first consider the case when (Z, g_Z) is $(S^2 \times S^2, g_{S^2 \times S^2})$. We can glue on an AF space at both points (n, n) and (s, s), but we must take the same AF space for both points. In this case, we impose an additional symmetry. There is an orientation-preserving involution of $S^2 \times S^2$ consisting of the product of antipodal maps. Since both AF spaces are the same, this involution obviously extends to an involution of $X_{a,b}$ which is an isometry of $g^{(0)}_{a,b}$, and which we will refer to as bilateral symmetry. As in the single bubble case, the toric action extends to an isometry of the approximate metric on the connect sum. We then have the following cases with toric invariance, diagonal symmetry, and bilateral symmetry:

- (iv) $3\# S^2 \times S^2$; the product metric on $S^2 \times S^2$ with Green's function $S^2 \times S^2$ metrics attached at two fixed points.
- (v) $S^2 \times S^2 \# 2\overline{\mathbb{CP}}^2 = \mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$; the product metric on $S^2 \times S^2$ with Burns metrics attached at two fixed points.

Next, we consider the case when (Z, g_Z) is $(\mathbb{CP}^2, g_{FS})$. Imposing trilateral symmetry (see Figure 2.1), allows us to attach the same AF space at all 3 fixed points. We then have the following cases with toric invariance, diagonal symmetry at each fixed point, and trilateral symmetry:

- (vi) $\mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$; the Fubini-Study metric with Burns metrics attached at all fixed points.

- (vii) $\mathbb{CP}^2\#3(S^2\times S^2) = 4\mathbb{CP}^2\#3\overline{\mathbb{CP}}^2$; the Fubini-Study metric with Green's function $S^2\times S^2$ metrics attached at all fixed points.

Next, we return to the case that (Z, g_Z) is $(S^2\times S^2, g_{S^2\times S^2})$. Imposing quadrilateral symmetry (see Figure 2.2), allows us to attach the same AF space at all 4 fixed points. We then have the following cases with toric invariance, diagonal symmetry at each fixed point, and quadrilateral symmetry:

- (viii) $S^2\times S^2\#4\overline{\mathbb{CP}}^2 = \mathbb{CP}^2\#5\overline{\mathbb{CP}}^2$; the product metric on $S^2\times S^2$ with Burns metrics attached at all fixed points.
- (ix) $5\#S^2\times S^2$ viewed as the product metric on $S^2\times S^2$ with Greens function $S^2\times S^2$ metrics attached at all fixed points.

For multiple bubbles in the non-orientable case, see Appendix B.

9.3. **Weight function.** For the weighted norms, we define the weight function on $X_{a,b}$ by

$$w = \begin{cases} a^{-1}b^{-1} & |z| \geq 1 \\ a^{-1}b^{-1}|z| & 1 \geq |z| \geq 2b \\ |x| & 2a^{-1} \geq |x| \geq 1 \\ 1 & 1 \geq |x|, \end{cases} \tag{9.16}$$

where for simplicity we have assumed that the x and z coordinates contain the unit spheres. We record the inequalities

$$1 \leq w \leq a^{-1}b^{-1}. \tag{9.17}$$

10. REFINED APPROXIMATE METRIC

Remark 10.1. We will now choose $\delta < 0$ satisyfing $-\epsilon < \delta < 0$, where ϵ was previously chosen (see Remark 7.6).

As pointed out above, the approximate metric defined in (9.9) is insufficient for our purposes, and needs to be refined. To define the new approximate metric, we replace g_N with $g_N^{(1)} = g_N + a^2b^2\tilde{H}_2(x)$, so that

$$g_N^{(1)} = g_0 + \eta_N(x) + a^2b^2\tilde{H}_2(x), \quad |x| \leq a^{-1}. \tag{10.1}$$

By Proposition 7.2, for a, b sufficiently small

$$\begin{aligned} |a^2b^2\tilde{H}_2(x)| &\lesssim a^2b^2|x|^2 \\ &\lesssim b^2, \end{aligned}$$

so that $g_N^{(1)}$ is indeed a Riemannian metric when $|x| < a^{-1}$.

Next, replace the compact metric g_Z with $g_Z^{(1)} = g_Z + a^2b^2\tilde{H}_{-2}(z)$, so that

$$g_Z^{(1)} = g_0 + \eta_Z(z) + a^2b^2\tilde{H}_{-2}(z), \quad |z| \geq b. \tag{10.2}$$

By Proposition 7.5,

$$\begin{aligned}|a^2b^2\tilde{H}_{-2}(z)| &\lesssim a^2b^2b^{-2}\\ &\lesssim a^2,\end{aligned}$$

hence $g_Z^{(1)}$ is a Riemannian metric for $|z| \geq b$.

Using these metrics, we then define the refined approximate metric $g_{a,b}^{(1)}$ on $X_{a,b}$ by

$$g_{a,b}^{(1)} = \begin{cases} a^{-2}b^{-2}(g_Z + a^2b^2\tilde{H}_{-2}(z)) & |z| \geq 2b,\\ g_N + a^2b^2\tilde{H}_2(x) & |x| \leq a^{-1}, \end{cases} \tag{10.3}$$

while in the damage zone $a^{-1} \leq |x| \leq 2a^{-1}$ the metric is given by

$$\begin{aligned} g_{a,b}^{(1)} = dx^2 &+ \phi(a|x|)\big\{\eta_N(x) + a^2b^2\tilde{H}_2(x)\big\}\\ &+ [1-\phi(a|x|)]\iota^*\big\{a^{-2}b^{-2}(\eta_Z(z) + a^2b^2\tilde{H}_{-2}(z))\big\}. \end{aligned} \tag{10.4}$$

Remark 10.2. From Propositions 7.2 and 7.5, it is clear that $g_{a,b}^{(1)}$ is invariant under the group action.

10.1. **Damage zone estimate.** We compute that

$$\begin{aligned} \iota^*\big\{a^{-2}b^{-2}(g_Z + a^2b^2\tilde{H}_{-2}(z))\big\} &= \iota^*\big\{a^{-2}b^{-2}(\delta_{ij} + (\eta_Z(z))_{ij} + a^2b^2\tilde{H}_{-2}(z)_{ij})dz^idz^j\big\}\\ &= \big(\eta_Z(abx)_{ij} + a^2b^2\tilde{H}_{-2}(abx)_{ij}\big)dx^idx^j. \end{aligned} \tag{10.5}$$

Consequently, in the damage zone, the metric is

$$\begin{aligned} g_{a,b}^{(1)} = dx^2 &+ \phi(a|x|)\big\{\eta_N(x) + a^2b^2\tilde{H}_2(x)\big\}\\ &+ [1-\phi(a|x|)]\big(\eta_Z(abx)_{ij} + a^2b^2\tilde{H}_{-2}(abx)_{ij}\big)dx^idx^j. \end{aligned} \tag{10.6}$$

We next use the the expansions

$$a^2b^2\tilde{H}_2(x)_{ij} = -a^2b^2\Big(\frac{1}{3}R_{ikjl}(z_0)x^kx^l + O^{(4)}(|x|^\epsilon)\Big), \tag{10.7}$$

$$\eta_Z(abx)_{ij} = -a^2b^2\frac{1}{3}R_{ikjl}(z_0)x^kx^l + a^4b^4O^{(4)}(|x|^4), \tag{10.8}$$

$$\begin{aligned} a^2b^2\tilde{H}_{-2}(abx)_{ij} &= a^2b^2\Big(H_{-2}(abx)_{ij} + (ab)^{-\epsilon}O(|x|^{-\epsilon})\Big)\\ &= -\frac{1}{3}R_{ikjl}(y_0)\frac{x^kx^l}{|x|^4} + 2A\frac{1}{|x|^2}\delta_{ij} + (ab)^{2-\epsilon}O^{(4)}(|x|^{-\epsilon}), \end{aligned} \tag{10.9}$$

and

$$\eta_N(x)_{ij} = -\frac{1}{3}R_{ikjl}(y_0)\frac{x^kx^l}{|x|^4} + 2A\frac{1}{|x|^2}\delta_{ij} + O^{(4)}(|x|^{-4+\epsilon}). \tag{10.10}$$

Using (10.7)-(10.10), we obtain in the damage zone:

$$\begin{aligned}(g^{(1)}_{a,b})_{ij} &= \delta_{ij} - a^2b^2\frac{1}{3}R_{ikjl}(z_0)x^kx^l - \frac{1}{3}R_{ikjl}(y_0)\frac{x^kx^l}{|x|^4} + 2A\frac{1}{|x|^2}\delta_{ij} \\ &\quad + a^2b^2O^{(4)}(|x|^{\epsilon}) + a^4b^4O^{(4)}(|x|^4) + (ab)^{2-\epsilon}O^{(4)}(|x|^{-\epsilon}) + O^{(4)}(|x|^{-4+\epsilon}).\end{aligned} \tag{10.11}$$

Proposition 10.3. *The size of the B^t-tensor of the refined approximate metric in the damage zone is given by*

$$|B^t(g^{(1)}_{a,b})|_{g^{(1)}_{a,b}} = O(b^2a^{6-\epsilon}) + O(a^4b^4) + O(a^6b^{2-\epsilon}) + O(a^{8-\epsilon}), \tag{10.12}$$

as $a, b \to 0$.

Proof. By (10.11),

$$g^{(1)}_{a,b} = g_0 + a^2b^2H_2 + H_{-2} + \mathcal{E},$$

where

$$\mathcal{E} = a^2b^2O^{(4)}(|x|^{\epsilon}) + a^4b^4O^{(4)}(|x|^4) + (ab)^{2-\epsilon}O^{(4)}(|x|^{-\epsilon}) + O^{(4)}(|x|^{-4+\epsilon}). \tag{10.13}$$

Let $\theta = g^{(1)}_{a,b} - g_0$; then using the expansion of the B^t-tensor in Proposition 3.8

$$\begin{aligned}B^t(g^{(1)}_{a,b}) &= B^t(g_0) + (B^t_0)'(\theta) + \mathcal{Q}_0(\theta) \\ &= (B^t_0)'(\theta) + \mathcal{Q}_0(\theta).\end{aligned} \tag{10.14}$$

By Lemma 7.3 and the fact that $(B^t_0)'$ is fourth order,

$$\begin{aligned}(B^t_0)'(\theta) &= a^2b^2(B^t_0)'(H_2) + (B^t_0)'(H_{-2}) + (B^t_0)'(\mathcal{E}) \\ &= (B^t_0)'(\mathcal{E}),\end{aligned}$$

hence

$$B^t(g^{(1)}_{a,b}) = (B^t_0)'(\mathcal{E}) + \mathcal{Q}_0(\theta). \tag{10.15}$$

If we estimate the norm of $B^t(g^{(1)}_{a,b})$ in the flat metric, then (10.12) follows from the formula for $\mathcal{E}$ and the result of Proposition 3.8. However, by (10.11) it is clear that the same estimate holds if we use the norm with respect to $g^{(1)}_{a,b}$, since for any symmetric $(0,2)$-tensor $T = T_{ij}$

$$\begin{aligned}|T|^2_{g^{(1)}_{a,b}} &= \{g^{(1)}_{a,b}\}^{ik}\{g^{(1)}_{a,b}\}^{j\ell}T_{ij}T_{k\ell} \\ &= \{\delta_{ik} + O(a^2+b^2)\}\{\delta_{j\ell} + O(a^2+b^2)\}^{j\ell}T_{ij}T_{k\ell} \\ &= \big(1 + O(a^2+b^2)\big)|T|^2_0.\end{aligned} \tag{10.16}$$

□

Next, on the asymptotically flat piece we have

Proposition 10.4. *On $\{|x| < a^{-1}\} \subset N$, the B^t-tensor satisfies*

$$B^t(g^{(1)}_{a,b}) = a^2b^2\lambda k_1^{(0)} - a^2b^2\mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2 + O(a^4b^4), \tag{10.17}$$

and

$$B^t(g^{(1)}_{a,b}) = a^2b^2\lambda k_1^{(0)} + a^2b^2 O(|x|^{\epsilon-4}) + O(a^4b^4), \tag{10.18}$$

as $a, b \to 0$ and $|x| \to \infty$.

Proof. The proof proceeds along the same lines as the proof of Proposition 10.3. On $\{|x| < a^{-1}\} \subset N$ we have

$$g^{(1)}_{a,b} = g_N + a^2b^2\tilde{H}_2. \tag{10.19}$$

Let $\theta = g^{(1)}_{a,b} - g_N = a^2b^2\tilde{H}_2$. Using the expansion of the B^t-tensor again,

$$\begin{aligned} B^t(g^{(1)}_{a,b}) &= B^t(g_N) + (B^t_{g_N})'\theta + \mathcal{Q}_N(\theta) \\ &= (B^t_{g_N})'\theta + \mathcal{Q}_N(\theta), \end{aligned} \tag{10.20}$$

since g_N is B^t-flat. From the formula for the linearized operator S in (3.11) and Proposition 7.2 it follows that

$$\begin{aligned} (B^t_{g_N})'\theta &= S_{g_N}\theta - \mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\theta} \\ &= a^2b^2S(\tilde{H}_2) - a^2b^2\mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2 \\ &= a^2b^2\lambda k_1^{(0)} - a^2b^2\mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2. \end{aligned} \tag{10.21}$$

Substituting this into (10.20) gives

$$B^t(g^{(1)}_{a,b}) = a^2b^2\lambda k_1^{(0)} - a^2b^2\mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2 + \mathcal{Q}_N(\theta). \tag{10.22}$$

By part (i) of Proposition 3.7,

$$|Q_N(a^2b^2\tilde{H}_2)|_{g_N} = O(a^4b^4), \tag{10.23}$$

and from our observations above the same estimate holds if we estimate with respect to the norm induced by $g^{(1)}_{a,b}$. Therefore, (10.17) follows from this estimate and (10.22).

To estimate the gauge-fixing term in (10.22), we first observe that $\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N} : S^2(T^*N) \to T^*N$ is a third order differential operator, while

$$\nabla^3_N = \partial^3 + \Gamma(g_N)\partial^2 + [\partial\Gamma(g_N) + \Gamma(g_N)^2]\partial + [\partial^2\Gamma(g_N) + \Gamma(g_N)\partial\Gamma(g_N)],$$

where ∇_N denotes the covariant derivative and $\Gamma(g_N)$ the Christoffel symbols in the g_N-metric. Since

$$\begin{aligned} \tilde{H}_2 &= H_2 + O^{(4)}(|x|^{\epsilon}), \\ \Gamma_N &= O^{(3)}(|x|^{-3}), \end{aligned}$$

it follows that

$$a^2b^2\mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2 = a^2b^2O(|x|^{\epsilon-4}),$$

and combining this with (10.23) and (10.22) we obtain (10.18). □

Next, on the compact piece, we have

Proposition 10.5. *On* $\{|z| \geq 2b\} \subset Z$*, we have*

$$|B^t(g^{(1)}_{a,b})|_{g^{(1)}_{a,b}} = a^6b^6O(|z|^{-\epsilon-4}). \tag{10.24}$$

as $a, b, |z| \to 0$.

Proof. Recall the metric $g^{(1)}_Z$ defined in (10.2):

$$g^{(1)}_Z = g_0 + \eta_Z(z) + a^2b^2\tilde{H}_{-2}(z), \quad |z| \geq 2b,$$

so that on the compact piece $\{|z| \geq 2b\} \cap Z$ the refined approximate metric $g^{(1)}_{a,b}$ is just a rescaling of $g^{(1)}_Z$:

$$g^{(1)}_{a,b} = a^{-2}b^{-2}g^{(1)}_Z. \tag{10.25}$$

We can then essentially repeat the arguments of the preceding propositions and write $g^{(1)}_Z = g_Z + \theta$, where $\theta = a^2b^2\tilde{H}_{-2}$, then expand B^t:

$$\begin{aligned} B^t(g^{(1)}_Z) &= B^t(g_Z) + (B^t_Z)'(\theta) + \mathcal{Q}_Z(\theta) \\ &= (B^t_Z)'(\theta) + \mathcal{Q}_Z(\theta), \end{aligned} \tag{10.26}$$

where as usual the subscript Z indicates that the tensor is with respect to the metric g_Z.

We first estimate the term involving the linearization of B^t, by comparing $(B^t_Z)'(\theta)$ and $(B^t_0)'(\theta)$, i.e., the linearized operator with respect to the flat metric acting on θ. Recall from Lemma 7.3 and Proposition 7.5 that

$$\begin{aligned} (B^t_0)'(\theta) &= a^2b^2(B^t_0)'(\tilde{H}_{-2}) \\ &= a^2b^2(B^t_0)'\big(H_{-2} + O(|z|^{-\epsilon})\big) \\ &= a^2b^2O(|z|^{-\epsilon-4}). \end{aligned} \tag{10.27}$$

Clearly, for any metric g the operators $(B^t)'$ and S have the same general form, as given in (7.6):

$$\begin{aligned} (B^t)'\theta = (g^{-2} + g * g^{-3}) * \nabla^4\theta + g * g^{-3} * Rm * \nabla^2\theta + g * g^{-3} * \nabla Rm * \nabla\theta \\ + (g^{-2} + g * g^{-3}) * (\nabla^2 Rm + Rm * Rm) * \theta. \end{aligned} \tag{10.28}$$

Using this, we can estimate the difference

$$[(B^t_Z)' - (B^t_0)'](\theta).$$

We will need the following estimates, which follow from (3.35), (9.1), and (9.2):

$$\begin{aligned} g_Z - g_0 &= O(b^2), \\ \nabla_Z^m T - \nabla_0^m T &= O(b) * \nabla_0^{m-1} T + O(1) * \{\nabla_0^{m-2} T + \cdots + T\}. \end{aligned} \tag{10.29}$$

Then by (10.28),

$$|[(B_Z^t)' - (B_0^t)'](\theta)| = a^2 b^2 O(|z|^{-4}). \tag{10.30}$$

Combining with (10.27), we obtain

$$|(B_Z^t)'(\theta)|_0 = a^2 b^2 O(|z|^{-\epsilon-4}). \tag{10.31}$$

It is easy to see that the same estimate holds if we estimate with respect to the norm induced by $g_Z^{(1)}$.

For the remainder term $\mathcal{Q}$ in (10.26) we use Proposition 3.8 to show

$$|\mathcal{Q}_Z(\theta)|_{g_Z} = a^4 b^4 O(|z|^{-8}), \tag{10.32}$$

with the same estimate in the $g_Z^{(1)}$-metric. Combining (10.31) and (10.32) gives

$$|B^t(g_Z^{(1)})|_{g_Z^{(1)}} = a^2 b^2 O(|z|^{-\epsilon-4}) + a^4 b^4 O(|z|^{-6}). \tag{10.33}$$

Since $|z| \geq b$, this implies

$$|B^t(g_Z^{(1)})|_{g_Z^{(1)}} = a^2 b^2 O(|z|^{-\epsilon-4}). \tag{10.34}$$

By the scaling properties of the B^t-tensor,

$$\begin{aligned} B^t(g_{a,b}^{(1)}) &= B^t(a^{-2} b^{-2} g_Z^{(1)}) \\ &= a^2 b^2 B^t(g_Z^{(1)}). \end{aligned}$$

Using (10.34) and (10.25) we conclude

$$\begin{aligned} |B^t(g_{a,b}^{(1)})|_{g_{a,b}^{(1)}} &= |a^2 b^2 B^t(g_Z^{(1)})|_{g_{a,b}^{(1)}} \\ &= a^2 b^2 |a^2 b^2 B^t(g_Z^{(1)})|_{g_Z^{(1)}} \\ &= a^4 b^4 |B^t(g_Z^{(1)})|_{g_Z^{(1)}} \\ &= a^6 b^6 O(|z|^{-\epsilon-4}). \end{aligned}$$

□

Finally, we have

Proposition 10.6. *Choosing $a = b$, we have*

$$\|B^t(g_{a,a}^{(1)}) - a^4 \lambda k_1^{(0)}\|_{C^{0,\alpha}_{\delta-4}(X_{a,a})} = O(a^{4+\delta-\epsilon}) \tag{10.35}$$

as $a \to 0$.

Proof. We begin estimating the leading term in the $C^{0,\alpha}_{\delta-4}$-norm. On the damage zone, $k_1^{(0)} = 0$, so by (10.12)

$$w^{4-\delta}|B^t(g^{(1)}_{a,a}) - a^4\lambda k_1^{(0)}| = O(a^{\delta-4}a^{8-\epsilon}) = O(a^{4+\delta-\epsilon}). \tag{10.36}$$

By (10.18), on the AF piece

$$\begin{aligned} w^{4-\delta}|B^t(g^{(1)}_{a,a}) - a^4\lambda k_1^{(0)}| &= |x|^{4-\delta}\{a^4 O(|x|^{\epsilon-4}) + O(a^8)\} \\ &= a^4 O(|x|^{\epsilon-\delta}) + a^8 O(|x|^{4-\delta}). \end{aligned} \tag{10.37}$$

Recall from Remark 10.1 that $-\epsilon < \delta < 0$; hence $\epsilon - \delta > 0$ and

$$w^{4-\delta}|B^t(g^{(1)}_{a,a}) - a^4\lambda k_1^{(0)}| = O(a^{4+\delta-\epsilon}). \tag{10.38}$$

On the compact piece, by (10.24)

$$\begin{aligned} w^{4-\delta}|B^t(g^{(1)}_{a,a}) - a^4\lambda k_1^{(0)}| &= a^{2\delta-8} O(|z|^{4-\delta})\{a^{12} O(|z|^{-\epsilon-4})\} \\ &= a^{2\delta+4} O(|z|^{-\delta-\epsilon}) \\ &= O(a^{4+\delta-\epsilon}), \end{aligned} \tag{10.39}$$

since $|z| \geq 2a$ and $-\epsilon < \delta < 0$ (see Remark 10.1).

For estimating the Hölder part of the weighted norm, one must use the formula (3.52) in the proof of Proposition 3.7. For example, the term with $h * \nabla^4 h$ is estimated like

$$\begin{aligned} & w^{4-\delta}(x_0)\frac{|(h * \nabla^4 h)(x_1) - (h * \nabla^4 h)(x_2)|}{|d(x_1,x_2)|^\alpha} \\ & \leq w^{-\delta}(x_0)|h(x_1)| \cdot w^{4-\delta}(x_0)\frac{|\nabla^4 h(x_1) - \nabla^4 h(x_2)|}{d(x_1,x_2)^\alpha} \\ & + w^{-\delta}(x_0)\frac{|h(x_1) - h(x_2)|}{|d(x_1,x_2)|^\alpha} \cdot w^{4-\delta}(x_0)|\nabla^4 h(x_2)|, \end{aligned} \tag{10.40}$$

and all other terms are estimated similarly, the complete computation is lengthy but straightforward, so is omitted. ☐

10.2. **The approximate cokernel.** In this subsection, we define tensors o_1, o_2, o_3 and k_1, k_2, k_3 which will be crucial in the Lyapunov-Schmidt reduction in Section 11.

Remark 10.7. It is clear that all of the tensors in this section may be chosen to be invariant under the group action, so we will do this automatically without mention in every case.

Recall from Section 5 we denoted the cokernel of the asymptotically flat manifold (N, g_N) by o_1, and it is given by

$$o_1 = \mathcal{K}\omega_1 + f g_N. \tag{10.41}$$

In Section 7 we defined a compactly supported symmetric $(0,2)$-tensor $k_1^{(0)} \in C_{\delta-4}$ which satisfies

$$\|k_1^{(0)}\|_{C_{\delta-4}} \leq C_1, \tag{10.42}$$

$$\int \langle k_1^{(0)}, o_1 \rangle_{g^{(0)}} \, dV_{g^{(0)}} = 1, \tag{10.43}$$

where C_1 is independent of a, b. Note that the quantities in (10.42) and (10.43) are all computed with respect to the "naive" approximate metric $g^{(0)}$ defined in Section 9. Since from now on we will be working in the refined approximate metric $g^{(1)}$ defined in Section 10, we will need to slightly rescale $k_1^{(0)}$ so that (10.42) and (10.43) hold with respect to $g^{(1)}$. To this end, define

$$k_1 = \Big\{ \int \langle k_1^{(0)}, o_1 \rangle_{g^{(1)}} \, dV_{g^{(1)}} \Big\}^{-1} k_1^{(0)}. \tag{10.44}$$

Then by (10.43) and (10.44),

$$\int \langle k_1, o_1 \rangle_{g^{(1)}} \, dV_{g^{(1)}} = 1. \tag{10.45}$$

Claim 10.8. *We have*

$$k_1^{(0)} = \big(1 + O(b^2)\big) k_1. \tag{10.46}$$

Proof. By (10.3), on the support of $k_1^{(0)}$ we have

$$\begin{aligned} g = g^{(1)} &= g_N + O(b^2) \\ &= g^{(0)} + O(b^2), \end{aligned} \tag{10.47}$$

hence the volume forms satisfy

$$dV_{g^{(1)}} = \big(1 + O(b^2)\big) dV_{g^{(0)}}. \tag{10.48}$$

Therefore,

$$\begin{aligned} \int \langle k_1^{(0)}, o_1 \rangle_{g^{(1)}} \, dV_{g^{(1)}} &= \int [g^{(1)}]^{ik} [g^{(1)}]^{j\ell} [k_1^{(0)}]_{ij} [o_1]_{k\ell} \, dV_{g^{(1)}} \\ &= \int \big\{ [g^{(0)}]^{ik} + O(b^2) \big\} \big\{ [g^{(0)}]^{j\ell} + O(b^2) \big\} [k_1^{(0)}]_{ij} [o_1]_{k\ell} \, \big(1 + O(b^2)\big) dV_{g^{(0)}} \\ &= \int \langle k_1^{(0)}, o_1 \rangle_{g^{(0)}} \, dV_{g^{(0)}} + O(b^2) \\ &= 1 + O(b^2). \end{aligned} \tag{10.49}$$

Substituting this into (10.44) gives (10.46). ☐

Remark 10.9. From now on, all metric-dependent quantities will be with respect to $g = g^{(1)}$. To simplify the notation, we will suppress the superscript.

Let o_3 denote the cokernel element on the compact manifold (Z, g_Z) given by scaling of the metric:

$$o_3 = (ab)^{-2+\delta} g_Z. \tag{10.50}$$

(The reason for the scale factor will become apparent in a moment). Fix a smooth positive cut-off function ϕ_3 supported in $Z \setminus \{z_0\}$ with

$$\int \phi_3 |g_Z|^2 \, dV = 1, \tag{10.51}$$

where we again emphasize that the volume form and inner product are with respect to $g = g^{(1)}$. Define

$$k_3 = (ab)^{2-\delta} \phi_3 g_Z. \tag{10.52}$$

We claim that there is a constant C_2 such that

$$\|k_3\|_{C_{\delta-4}} \le C_2. \tag{10.53}$$

To see this, first recall that by (10.3), (10.3), on the support of ϕ_3 we have

$$\begin{aligned} [g^{(1)}]_{ij} &= (ab)^{-2} \big[(g_Z)_{ij} + O(b^2) \big], \\ [g^{(1)}]^{ij} &= (ab)^{2} \big[(g_Z)^{ij} + O(b^2) \big]. \end{aligned}$$

Also, on the support of ϕ_3 the weight $w(x) \approx (ab)^{-2}$. It follows that

$$\begin{aligned} \sup |k_3|_{g^{(1)}} w^{4-\delta} &= c_2 (ab)^{2-\delta} \sup |\phi_3| |g_Z|_{g^{(1)}} w^{4-\delta} \\ &= c_2 (ab)^{2-\delta} \sup |\phi_3| \big\{ [g^{(1)}]^{ik} [g^{(1)}]^{j\ell} (g_Z)_{ij} (g_Z)_{k\ell} \big\}^{1/2} w^{4-\delta} \\ &= c_2 (ab)^{2-\delta} \sup \Big\{ |\phi_3| \big\{ (ab)^4 \big[(g_Z)^{ik} + O(b^2) \big] \\ &\qquad \times \big[(g_Z)^{j\ell} + O(b^2) \big] (g_Z)_{ij} (g_Z)_{k\ell} \big\}^{1/2} w^{4-\delta} \Big\} \\ &\le C_2. \end{aligned}$$

This estimate clarifies the choice of scaling in the definitions of k_3 and o_3: the scale factor $(ab)^{2-\delta}$ in (10.52) is necessary to get the bound (10.53), while the factor $(ab)^{-2+\delta}$ in the definition of o_3 was chosen to give the normalization

$$\begin{aligned} \int \langle k_3, o_3 \rangle \, dV &= \int \langle (ab)^{2-\delta} \phi_3 g_Z, (ab)^{-2+\delta} g_Z \rangle \, dV \\ &= \int \phi_3 |g_Z|^2 \, dV = 1. \end{aligned} \tag{10.54}$$

Next, denote

$$o_2 = g = g^{(1)}. \tag{10.55}$$

We claim that there is a tensor k_2, compactly supported in N, which satisfies the normalization

$$\int \langle k_2, o_2 \rangle \, dV = \int \mathrm{tr}_g k_2 \, dV = 1 \tag{10.56}$$

and the orthogonality condition

$$\int \langle k_2, o_1 \rangle \, dV = 0. \tag{10.57}$$

(Note that the integral in (10.57) makes sense, since k_2 is compactly supported in N, even though o_1 is not globally defined.)

To see that such a tensor exists, just take two smooth, positive cut-off functions η_1, η_2 with compact support in N and let

$$k_2 = (c_1\eta_1 + c_2\eta_2)o_2 = (c_1\eta_1 + c_2\eta_2)g,$$

where c_1 and c_2 are constants to be determined. Then

$$\begin{aligned} \int \langle k_2, o_1 \rangle \, dV &= c_1 \int \eta_1 (\mathrm{tr}_g o_1) \, dV + c_2 \int \eta_2 (\mathrm{tr}_g o_1) \, dV, \\ \int \langle k_2, o_2 \rangle \, dV &= 4c_1 \int \eta_1 \, dV + 4c_2 \int \eta_2 \, dV. \end{aligned} \tag{10.58}$$

By (10.47), on the support of k_2

$$g = g^{(1)} = g_N + O(b^2).$$

Therefore,

$$\begin{aligned} \mathrm{tr}_g o_1 &= [g^{(1)}]^{k\ell} (o_1)_{k\ell} \\ &= [g_N + O(b^2)]^{k\ell} [\mathcal{K}\omega_1 + f g_N]_{k\ell} \\ &= 4f + O(b^2). \end{aligned}$$

Therefore, we can estimate the integrals in (10.58) by

$$\begin{aligned} \int \langle k_2, o_1 \rangle \, dV &= c_1 \Big\{ 4 \int \eta_1 f \, dV + O(b^2) \Big\} + c_2 \Big\{ 4 \int \eta_2 f \, dV + O(b^2) \Big\}, \\ \int \langle k_2, o_2 \rangle \, dV &= c_1 \Big\{ 4 \int \eta_1 \, dV \Big\} + c_2 \Big\{ 4 \int \eta_2 \, dV \Big\}. \end{aligned} \tag{10.59}$$

Consequently, once a, b are small enough it is possible to choose the cut-off functions η_1, η_2 and the constants c_1, c_2 so that (10.56) and (10.57) hold.

11. LYAPUNOV-SCHMIDT REDUCTION

In this section, we perform the main reduction of the problem from an infinite-dimensional problem to a finite-dimensional problem.

Remark 11.1. Since we are carrying out an equivariant gluing construction, from now on all operators are understood to act on sections of the relevant bundle which are invariant under the group actions described above.

11.1. **The modified nonlinear map.** Let

$$\mathcal{D} = \Big\{ h \in C^{4,\alpha}_{\delta}(X_{a,b}) \ : \ \int \langle h, k_1 \rangle = 0, \ \int \langle h, k_3 \rangle = 0 \ \Big\}. \tag{11.1}$$

Define the mapping $H^t : \mathbb{R} \times \mathbb{R} \times \mathcal{D} \to C^{0,\alpha}_{\delta-4}(X_{a,b})$ by

$$H^t(\lambda_1, \lambda_2, \theta) = P_{g^{(1)}}(\theta) - \lambda_1 k_1 - \lambda_2 k_2. \tag{11.2}$$

Let $(H^t)' : \mathbb{R} \times \mathbb{R} \times \mathcal{D} \to C^{0,\alpha}_{\delta-4}$ denote the linearization of H at $(0,0,0)$:

$$(H^t)'(\lambda_1, \lambda_2, h) = \frac{d}{ds} H^t(s\lambda_1, s\lambda_2, sh)\Big|_{s=0}. \tag{11.3}$$

Then $(H^t)'$ is given by

$$(H^t)'(\lambda_1, \lambda_2, h) = S(h) - \lambda_1 k_1 - \lambda_2 k_2, \tag{11.4}$$

where $S = S^t$ is the linearization of P at $g^{(1)}$.

Proposition 11.2. *For a, b sufficiently small, the map $(H^t)' : \mathbb{R} \times \mathbb{R} \times \mathcal{D} \to C^{0,\alpha}_{\delta-4}$ is uniformly injective: i.e., there is a constant $\delta_0 > 0$ which is independent of a, b such that*

$$\|(H^t)'(\lambda_1, \lambda_2, h)\|_{C^{\alpha}_{\delta-4}} \geq \delta_0 \big(|\lambda_1| + |\lambda_2| + \|h\|_{C^{4,\alpha}_{\delta}} \big). \tag{11.5}$$

Proof. We argue via contradiction: if (11.5) does not hold, then there is a sequence $(\lambda_1^i, \lambda_2^i, h_i) \in \mathcal{D}$ with

$$\begin{aligned} &|\lambda_1^i| + |\lambda_2^i| + \|h_i\|_{C^{4,\alpha}_{\delta}} = 1 \ \ \forall i, \\ &\epsilon_i \equiv (H^t)'(\lambda_1^i, \lambda_2^i, h_i) \to 0 \ \text{ in } C^{0,\alpha}_{\delta-4}. \end{aligned} \tag{11.6}$$

If we pair ϵ_i with ηo_1 and integrate, where η is a cut-off function with

$$\eta(x) = \begin{cases} 1 & |x| \leq a^{-1} \\ 0 & |x| > 2a^{-1}, \end{cases} \tag{11.7}$$

then

$$\int \langle \epsilon_i, \eta o_1 \rangle \ dV = \int_B \langle S h_i, \eta o_1 \rangle \ dV - \lambda_1^i \int \langle k_1, \eta o_1 \rangle \ dV - \lambda_2^i \int \langle k_2, \eta o_1 \rangle \ dV. \tag{11.8}$$

Since $\eta \equiv 1$ on the support of k_1 and k_2, by the normalization (10.42) and the orthogonality condition (10.57) we can rewrite this as

$$\lambda_1^i = - \int \langle \epsilon_i, \eta o_1 \rangle \ dV + \int \langle S h_i, \eta o_1 \rangle \ dV. \tag{11.9}$$

For the first term on the right-hand side, note that

$$\|\epsilon_i\|_{C^{0,\alpha}_{\delta-4}} \geq \sup\{ |\epsilon_i| w^{4-\delta} \}, \tag{11.10}$$

where w is the weight function. According to (9.16), on the support of η the weight function is $w(x) = |x|$ (for $|x|$ large). Also, by Theorem 6.1 the cokernel o_1 satisfies

$$|o_1| \leq C |x|^{-2}.$$

Therefore,

$$
\begin{aligned}
\Big|\int \langle \epsilon_i, \eta o_1\rangle \, dV\Big| &\leq \int |\epsilon_i||\eta o_1| \, dV \\
&\leq C\|\epsilon_i\|_{C^{0,\alpha}_{\delta-4}} \int_{1<<|x|\leq a^{-1}/2} |x|^{\delta-4}|x|^{-2} \, dV \\
&\leq C\|\epsilon_i\|_{C^{0,\alpha}_{\delta-4}} \to 0
\end{aligned} \tag{11.11}
$$

as $i \to \infty$.

For the second term on the right-hand side of (11.9) we integrate by parts, using the fact that S is self-adjoint:

$$
\int \langle Sh_i, \eta o_1\rangle \, dV = \int \langle h_i, S(\eta o_1)\rangle \, dV. \tag{11.12}
$$

Using the formula for S in (7.6) and the Leibniz rule, write

(11.13)

$$
\begin{aligned}
S(\eta o_1) &= (g^{-2} + g * g^{-3}) * \nabla^4(\eta o_1) + g * g^{-3} * Rm * \nabla^2(\eta o_1) \\
&\quad + g * g^{-3} * \nabla Rm * \nabla(\eta o_1) + (g^{-2} + g * g^{-3}) * (\nabla^2 Rm + Rm * Rm) * (\eta o_1) \\
&= \eta S o_1 + (g^{-2} + g * g^{-3}) * \sum_{j\geq 1}^{4} \nabla^{4-j} o_1 * \nabla^j \eta \\
&\qquad + g * g^{-3} * Rm * \sum_{j\geq 1}^{2} \nabla^{2-j} o_1 * \nabla^j \eta + g * g^{-3} * \nabla Rm * o_1 * \nabla \eta.
\end{aligned}
$$

By (5.45) and Theorem 6.1, on the support of $|\nabla \eta|$

$$
\begin{aligned}
|\nabla^m o_1| &= O(a^{m+2}), \\
|\nabla^m \eta| &= O(a^m), \\
|\nabla^m Rm| &= O(a^{m+4}).
\end{aligned}
$$

Therefore, from (11.13) we have

$$
S(\eta o_1) = \eta S o_1 + \{\text{Error}\}, \tag{11.14}
$$

where the error is supported on $\{a^{-1}/4 \leq |x| \leq a^{-1}/2\}$ and satisfies

$$
|\{\text{Error}\}| = O(a^6). \tag{11.15}
$$

It follows from (11.12)

$$
\int \langle Sh_i, \eta o_1\rangle \, dV = \int \langle h_i, \eta S(o_1)\rangle \, dV + \int \langle h_i, \{\text{Error}\}\rangle \, dV. \tag{11.16}
$$

Since $\|h_i\|_{C^{4,\alpha}_{\delta}} \leq 1$,

$$
|h_i| \leq C w^{\delta}, \tag{11.17}
$$

hence on the support of Error

$$|h_i| \leq C a^{-\delta}, \tag{11.18}$$

hence by (11.15)

$$\Big| \int_{\{a^{-1}/4 \leq |x| \leq a^{-1}/2\}} \langle h_i, \{\text{Error}\} \rangle \, dV \Big| \leq C a^{2-\delta}. \tag{11.19}$$

Therefore,

$$\int \langle S h_i, \eta o_1 \rangle \, dV = \int \langle h_i, \eta S o_1 \rangle \, dV + O(a^{2-\delta}). \tag{11.20}$$

Let S_N denote the linearized operator with respect to the metric g_N. Then $S_N o_1 = 0$, hence

$$\begin{aligned} S o_1 &= (S - S_N) o_1 + S_N o_1 \\ &= (S - S_N) o_1. \end{aligned} \tag{11.21}$$

Using (7.6) with (10.47), we can estimate

$$\begin{aligned} |(S - S_N) o_1| &\lesssim b^2 |\nabla^4 o_1| + b^2 a |\nabla^3 o_1| + a^2 b^2 |\nabla^2 o_1| + a^3 b^2 |\nabla o_1| \\ &\quad + a^4 b^2 |o_1| + b^2 |Rm| |\nabla^2 o_1| + b^2 |\nabla Rm| |\nabla o_1| + b^2 \big(|\nabla^2 Rm| + |Rm|^2 \big) |o_1|. \end{aligned} \tag{11.22}$$

Therefore, by (11.18),

$$\begin{aligned} \Big| \int \langle h_i, \eta S o_1 \rangle \, dV \Big| &= \Big| \int \langle h_i, \eta (S - S_N) o_1 \rangle \, dV \Big| \\ &\leq C \int w^{\delta} |(S - S_N) o_1| \, dV \\ &\leq C b^2 a^{2-\delta}. \end{aligned} \tag{11.23}$$

Combining the above, we conclude

$$\int \langle S h_i, \eta o_1 \rangle \, dV = O(a^{2-\delta}), \tag{11.24}$$

hence by (11.9),

$$\lambda_1^i \to 0 \ \text{ as } i \to \infty. \tag{11.25}$$

Next, pair ϵ_i with ηo_2 and integrate:

$$\int \langle \epsilon_i, \eta o_2 \rangle \, dV = \int \langle S h_i, \eta o_2 \rangle \, dV - \lambda_1^i \int \langle k_1, \eta o_2 \rangle \, dV - \lambda_2^i \int \langle k_2, \eta o_2 \rangle \, dV. \tag{11.26}$$

By the normalization (10.56), we can rewrite this as

$$\lambda_2^i = - \int \langle \epsilon_i, \eta o_2 \rangle \, dV + \int \langle S h_i, \eta o_2 \rangle \, dV - \lambda_1^i \int \langle k_1, \eta o_2 \rangle \, dV \tag{11.27}$$

As in (11.11), we can estimate the first integral on the right as

$$\begin{aligned}\Big| \int \langle \epsilon_i, \eta o_2 \rangle \, dV \Big| &\leq \int |\epsilon_i||\eta o_2| \, dV \\ &\leq C \|\epsilon_i\|_{C^{0,\alpha}_{\delta-4}} \int_{1<|x|\leq a^{-1}/2} |x|^{\delta-4} \, dV \\ &\leq C_\delta \|\epsilon_i\|_{C^{0,\alpha}_{\delta-4}} \quad (\text{since } \delta < 0),\end{aligned} \tag{11.28}$$

which limits to 0 as $i \to \infty$. The second term on the right we estimate as we did above; namely,

$$\int \langle Sh_i, \eta o_2 \rangle \, dV = \int \langle h_i, S(\eta o_2) \rangle \, dV. \tag{11.29}$$

Using the fact that $\nabla o_2 = 0$, we can estimate as in (11.13):

$$\begin{aligned} S(\eta o_2) &= (g^{-2} + g * g^{-3}) * \nabla^4(\eta o_2) + g * g^{-3} * Rm * \nabla^2(\eta o_2) \\ &\quad + g * g^{-3} * \nabla Rm * \nabla(\eta o_2) + (g^{-2} + g * g^{-3}) * (\nabla^2 Rm + Rm * Rm) * (\eta o_2) \\ &= \eta S o_2 + (g^{-2} + g * g^{-3}) * \nabla^4 \eta * o_2 + g * g^{-3} * Rm * \nabla^2 \eta * o_2 \\ &\qquad + g * g^{-3} * \nabla Rm * o_2 * \nabla \eta. \end{aligned}$$

Since $So_2 = Sg = 0$,

$$\begin{aligned} S(\eta o_2) &= \eta S o_2 + \{\text{Error}\}, \\ &= \{\text{Error}\}, \end{aligned} \tag{11.30}$$

where the error is supported on $\{a^{-1}/4 \leq |x| \leq a^{-1}/2\}$ and satisfies

$$|\{\text{Error}\}| = O(a^4). \tag{11.31}$$

Using (11.18), we can therefore estimate

$$\Big| \int_{\{a^{-1}/4 \leq |x| \leq a^{-1}/2\}} \langle h_i, \{\text{Error}\} \rangle \, dV \Big| \leq C a^{-\delta}.$$

Hence,

$$\int \langle Sh_i, \eta o_2 \rangle \, dV = O(a^{-\delta}). \tag{11.32}$$

For the last term in (11.27), we use the fact that $\lambda_1^i \to 0$, and that k_1 is compactly supported:

$$\begin{aligned} \Big| \lambda_1^i \int \langle k_1, \eta o_2 \rangle \, dV \Big| &\leq |\lambda_1^i| \int \eta |k_1||o_2| \, dV \\ &\leq C |\lambda_1^i| \int_{\text{supp}\{k_1\}} dV \\ &\leq C |\lambda_1^i| \to 0, \end{aligned}$$

as $i \to 0$. Combining with (11.32), (11.28), and (11.27), we see that

$$\lambda_2^i \to 0 \tag{11.33}$$

as $i \to \infty$.

Consequently, by (11.6) we now know

$$\|h_i\|_{C^{4,\alpha}_{\delta}} \to 1, \tag{11.34}$$

$$\|Sh_i\|_{C^{0,\alpha}_{\delta-4}} \to 0, \tag{11.35}$$

as $i \to \infty$.

The remainder of the proof is a standard "blow-up" argument, which we only briefly outline. Let (a_i, b_i) be a sequence of gluing parameters with $(a_i, b_i) \to (0, 0)$ as $i \to \infty$, and let $p_i \in X_{a_i,b_i}$ a sequence of points at which the supremum in (11.34) is attained. We have the three possibilities:

- (1) $p_i \to p \in N$. In this case, standard elliptic estimates produce a nontrivial solution of the limiting equation $Sh_\infty = 0$ on (N, g_N) with $h \in C^{4,\alpha}_{\delta}$. By Theorem 5.1, $h_\infty = c \cdot o_1$ for some $c \in \mathbb{R}$. Since

$$\int \langle h_i, k_1 \rangle dV_{(a_i,b_i)} = 0, \tag{11.36}$$

  k_1 has compact support on N, and $h_i \leq C w^\delta$, the integrand is bounded. Therefore,

$$\int \langle h_\infty, k_1 \rangle dV_{g_N} = 0, \tag{11.37}$$

  which implies that $c = 0$, a contradiction.
- (2) $p_i \to p \in Z \setminus \{z_0\}$. In this case, define $\tilde{h}_i = (ab)^{2+\delta} h_i$. It is easy to see that this scaling preserves the $C^{4,\alpha}_{\delta}$ norm, with respect to the metric $\tilde{g}^{(1)}_{a,b} = (ab)^2 g^{(1)}_{a,b}$. Standard elliptic estimates produce a nontrivial solution of the limiting equation $S\tilde{h}_\infty = 0$ on (Z, g_Z) with $h \in C^{4,\alpha}_{\delta}$. By Theorem 4.1, $h_\infty = c \cdot g_Z$ for some $c \in \mathbb{R}$. Since

$$\int \langle h_i, k_3 \rangle dV_{(a_i,b_i)} = 0, \tag{11.38}$$

  scaling shows that

$$\int \langle \tilde{h}_i, \phi_3 g_Z \rangle_{\tilde{g}} dV_{\tilde{g}} = (ab)^{2\delta} \cdot 0 = 0. \tag{11.39}$$

  Since ϕ_3 has compact support on $Z \setminus \{z_0\}$, and $\tilde{h}_i \leq C \tilde{w}^\delta = C(abw)^\delta$, the integrand is bounded, which implies that

$$\int \langle \tilde{h}_\infty, k_3 \rangle dV_{g_Z} = 0, \tag{11.40}$$

  which implies that $c = 0$, a contradiction.
- (3) If neither of the above cases happen, then there are two possibilities: a subsequence can approach the damage zone from the AF side, or from the

compact side. We give the argument in the former case, the proof of the latter case is similar. Fix a point $O \in N$ and let

$$d_i = \mathrm{dist}_{g^{(1)}_{a_i,b_i}}(O, p_i).$$

Clearly, $d_i \to \infty$ as $i \to \infty$ (otherwise we are in case (1) above). For $i >> 1$ we can view the sequence $\{p_i\} \subset N_i = N \cap A_i$, where A_i is the annulus $\{R_0 < |x| < 2a_i^{-1}\}$ and N_i is equipped with the metric $g_i = g^{(1)}_{a_i,b_i}$. Let $\psi_i : A_i \to N$ denote dilation,

$$\psi_i : x \mapsto d_i x,$$

and define

$$\tilde{g}_i = d_i^{-2} \psi_i^* g_i \Big|_{\{m_i d_i \leq |x| \leq M_i d_i\}}, \tag{11.41}$$

where $m_i \to 0$ and $M_i \to \infty$ are chosen so that the annulus $\{m_i d_i \leq |x| \leq M_i d_i\} \subset N_i$. Denote the dilated coordinates by $\tilde{x}^i$; then $\tilde{g}_i$ is defined on the annulus $\{m_i \leq |\tilde{x}^i| \leq M_i\}$. Finally, define

$$\tilde{h}_i = d_i^{-2+\delta} \psi_i^* h_i,$$

which preserves the $C^{4,\alpha}_{\delta}$-norm. Taking the limit as $i \to \infty$ we have $\tilde{g}_i \to ds^2$, the flat metric on $\mathbb{R}^4 \setminus \{0\}$, $\tilde{h}_i \to h_\infty$, where h_∞ satisfies

$$\begin{aligned} S_0 h_\infty &= 0 \ \text{ on } \mathbb{R}^4 \setminus \{0\}, \\ h_\infty &\in C^{4,\alpha}_{\delta}(\mathbb{R}^4 \setminus \{0\}), \end{aligned} \tag{11.42}$$

and S_0 is the linearized operator with respect to the flat metric (see (3.13)). Note the weight function in the limit is given by $w = |x|$. Since $-1 < \delta < 0$, δ is not an indicial root by Proposition 4.4. This implies that $S_0 : C^{4,\alpha}_{\delta} \to C^{0,\alpha}_{\delta-4}$ is an isomorphism (see [Bar86, LM85]), so $h_\infty \equiv 0$, which is a contradiction.

This contradiction argument finishes the proof of Proposition 11.2. □

We next quote without proof the following standard implicit function theorem:

Lemma 11.3. *Let $H : E \to F$ be a smooth map between Banach spaces. Define $Q = H - H(0) - H'(0)$. Assume that there are positive constants C_1, s_0, C_2 so that the following are satisfied:*

- (1) *The nonlinear term Q satisfies*

$$\|Q(x) - Q(y)\|_F \leq C_1(\|x\|_E + \|y\|_E)\|x - y\|_E \tag{11.43}$$

  for every $x, y \in B_E(0, s_0)$.
- (2) *The linearized operator at 0, $H'(0) : E \to F$ is an isomorphism with inverse bounded by C_2.*

If

$$s < \min\left(s_0, \frac{1}{2C_1C_2}\right) \tag{11.44}$$

and

$$\|H(0)\|_F < \frac{s}{2C_2}, \tag{11.45}$$

Then there is a unique solution $x \in B_E(0,s)$ *of the equation* $H(x) = 0$.

We end this section with the following existence theorem:

Theorem 11.4. *Let* $a = b$. *Then for all* a *sufficiently small, there exist constants* $\lambda_1, \lambda_2 \in \mathbb{R}$ *and* $\theta \in \mathcal{D}$ *satisyfying*

$$\|\theta\|_{C^{4,\alpha}_\delta} < Ca^{4+\delta-\epsilon} \tag{11.46}$$

so that

$$P_{g^{(1)}}(\theta) = \lambda_1 k_1 + \lambda_2 k_2. \tag{11.47}$$

Proof. We denote the refined approximate metric by $g^{(1)} = g^{(1)}_a$, or by g if the context is clear.

We will find a zero of H, so we need to verify the assumptions in Lemma 11.3 with $E = \mathbb{R} \times \mathbb{R} \times \mathcal{D}$ and $F = C^{0,\alpha}_\delta$, beginning with (1):

Lemma 11.5. *The quadratic estimate* (11.43) *holds for* $H : \mathbb{R} \times \mathbb{R} \times \mathcal{D} \to C^{0,\alpha}_\delta$.

Proof. This follows from Proposition 3.7, once we verify the assumptions (3.29) (the assumptions (3.28) clearly hold). We need to verify the estimate on each of the three regions: the asymptotically flat piece, the damage zone, and the compact piece. Recall that the weight is given by (9.16).

On the asymptotically flat piece, i.e., for $|x| \leq a^{-1}$,

$$g^{(1)} = g_N + a^4 \tilde{H}_2.$$

Let $h_1 = g^{(1)} - g_N = a^4 \tilde{H}_2$, then using the formula (3.36) we have

$$Rm_{g^{(1)}} = Rm_{g_N} + (g^{(1)})^{-1} * \nabla^2_N h_1 + (g^{(1)})^{-2} * \nabla_N h_1 * \nabla h_1.$$

By Proposition 7.2 and the fact that g_N is asymptotically flat of order 2,

$$\begin{aligned} Rm_{g^{(1)}} &= O(|x|^{-4}) + O(a^4) + O(a^8|x|^2) \\ &= O(|x|^{-4}) + O(a^4). \end{aligned} \tag{11.48}$$

Since $w(x) = |x|$ for $|x| >> 1$, it follows that

$$w(x)^2 |Rm_{g^{(1)}}| \leq C_0. \tag{11.49}$$

Similarly, using (3.37)

$$\nabla_{g^{(1)}} Rm_{g^{(1)}} = O(|x|^{-5}) + O(a^4|x|^{-3}) + O(a^4|x|^{\epsilon-3}) + O(a^8|x|) + O(a^{12}|x|^3), \tag{11.50}$$

hence

$$w(x)^3 |\nabla_{g^{(1)}} Rm_{g^{(1)}}| \leq C_1. \tag{11.51}$$

Finally, (3.38)

$$\begin{aligned}\nabla^2_{g^{(1)}} Rm_{g^{(1)}} &= O(|x|^{-6}) + O(a^4|x|^{-4}) + O(a^8|x|^{-2}) + O(a^4|x|^{-4+\epsilon}) \\ &\quad + O(a^8|x|^{\epsilon-2}) + O(a^8|x|^{2\epsilon-4}) + O(a^{12}|x|^2) + O(a^{16}|x|^4).\end{aligned} \tag{11.52}$$

Therefore,

$$w(x)^4 |\nabla^2_{g^{(1)}} Rm_{g^{(1)}}| \leq C_2. \tag{11.53}$$

The estimates for the other regions are verified in a similar manner, so we omit the details. □

It remains to show that

$$H' : \mathbb{R} \times \mathbb{R} \times \mathcal{D} \to C^{0,\alpha}_{\delta-4} \tag{11.54}$$

is an isomorphism with bounded inverse. This will follow once we prove surjectivity; the bound on the inverse will then follow immediately from Proposition 11.2.

In the following, let us view H' as a map

$$H' : \mathbb{R} \times \mathbb{R} \times C^{4,\alpha}_{\delta} \to C^{0,\alpha}_{\delta-4}. \tag{11.55}$$

Then the formal adjoint of H' maps from

$$(H')^* : C^{4,\alpha}_{-\delta} \to \mathbb{R} \times \mathbb{R} \times C^{0,\alpha}_{-\delta-4} \tag{11.56}$$

and is given by

$$(H')^*(h) = \Big(\int \langle h, k_1 \rangle dV, \int \langle h, k_2 \rangle dV, Sh \Big) \tag{11.57}$$

since S is self-adjoint (the duals of Hölder spaces are not Hölder spaces, but this slight abuse of notation should not cause confusion). We claim that for a sufficiently small, $Ker((H')^*) = 0$. To see this argue by contradiction: let h_i be a sequence of kernel elements corresponding to a sequence $a_i \to 0$ as $i \to \infty$. Normalize h_i so that $\|h_i\|_{C^{4,\alpha}_{-\delta}} = 1$. We then have a sequence h_i satisfying

$$Sh_i = 0, \quad \int \langle h, k_1 \rangle dV = 0, \quad \int \langle h, k_2 \rangle dV = 0, \tag{11.58}$$

$$\|h_i\|_{C^{4,\alpha}_{-\delta}} = 1. \tag{11.59}$$

The limiting argument in the proof of Proposition 11.2 is then modified as follows. Let p_i be a sequence of points in X_{a_i,b_i} for a sequence $a_i \to 0$ as $i \to \infty$ at which the supremum in the norm (11.59) is attained. We have the three possibilities.

- (1) $p_i \to p \in N$. In this case, standard elliptic estimates produce a nontrivial solution of the limiting equation $Sh_\infty = 0$ on (N, g_N) with $h \in C^{4,\alpha}_{-\delta}$. By Theorem 5.1, $h_\infty = c_1 \cdot o_1 + c_2 g_N$ for some $c_1, c_2 \in \mathbb{R}$. Since

$$\int \langle h_i, k_1 \rangle dV_{g^{(1)}_{a_i}} = 0, \quad \int \langle h_i, k_2 \rangle dV_{g^{(1)}_{a_i}} = 0, \tag{11.60}$$

and k_1, k_2 both have compact support on N, and $h_i \leq Cw^{-\delta}$, the integrand is bounded, which implies that

$$\int \langle h_\infty, k_1 \rangle dV_{g_N} = 0, \quad \int \langle h_\infty, k_2 \rangle dV_{g_N} = 0, \tag{11.61}$$

which implies that $c_1 = c_2 = 0$, a contradiction.

- (2) $p_i \to p \in Z \setminus \{z_0\}$. In this case, defined $\tilde{h}_i = (a_i)^{4+2\delta} h_i$. It is easy to see that this scaling preserves the $C^{4,\alpha}_\delta$ norm, with respect to the metric $\tilde{g}^{(1)}_{a_i} = a^4 g^{(1)}_{a_i}$. Standard elliptic estimates produce a nontrivial solution of the limiting equation $S\tilde{h}_\infty = 0$ on (Z, g_Z) with $h \in C^{4,\alpha}_{-\delta}$. By Theorem 4.1, $h_\infty = 0$ which is a contradiction.
- (3) If neither of the above cases happen, then as above one can rescale both the metric and h_i to find a solution $h_\infty \in C^{4,\alpha}_{-\delta}$ of the equation $Sh_\infty = 0$ on $\mathbb{R}^4 \setminus \{0\}$ with weight function $w = r$. Since $0 < -\delta < 1$, δ is not an indicial root so $S : C^{4,\alpha}_\delta \to C^{0,\alpha}_{\delta-4}$ is an isomorphism, therefore $h_\infty = 0$.

This contradiction proves that $Ker((H')^*) = \{0\}$, and by standard Fredholm Theory, we conclude that

$$H' : \mathbb{R} \times \mathbb{R} \times C^{4,\alpha}_\delta \to C^{0,\alpha}_{\delta-4}.$$

is surjective.

Claim 11.6. *For a sufficiently small, the dimension of the kernel of* $H' : \mathbb{R} \times \mathbb{R} \times C^{4,\alpha}_\delta \to C^{0,\alpha}_{\delta-4}$ *is at least 2.*

Proof. To see this, we claim that k_1 and k_2 are not in the image of S. If, for example $Sh_i = k_1$, then a limiting argument as above would produce a solution of $S(h_\infty) = k_1$ on (N, g_N), which is a contradiction. Similarly, if $Sh_i = k_2$, the same argument yields a contradiction. We have found 2 linearly independent elements not in the image of S; by Fredholm theory the cokernel of S must be at least two-dimensional. Since S is a self-adjoint operator, we must have $\dim(Ker(S)) \geq 2$. Obviously $\{0\} \times \{0\} \times Ker(S) \subset Ker(H')$, so the claim follows. □

To finish, by standard L^2-decomposition

$$L^2 \cap C^{4,\alpha}_\delta = \mathrm{span}\{k_1, k_3\} \oplus \mathcal{D}, \tag{11.62}$$

where $\mathcal{D} = C^{4,\alpha}_\delta \cap (\mathrm{span}\{k_1, k_3\})^\perp$. Let $h_1, h_2, \ldots h_j$ be a basis for $Ker(H')$, where $j = \dim(Ker(H'))$. Then we can write

$$h_i = c_{i1}k_1 + c_{i2}k_3 + m_i, \tag{11.63}$$

where $m_i \in \mathcal{D}$. If $j > 2$, then obviously we can take a nontrivial linear combination to obtain

$$\sum_i c_i h_i = \sum_i c_i m_i \tag{11.64}$$

for some constants c_i. The left hand side is in the kernel of H', but Proposition 11.2 shows that the left hand side cannot be, which is a contradiction. Consequently, from Claim 11.6 we conclude that $\dim(Ker(H')) = 2$. So we have the equations

$$h_1 = c_{11}k_1 + c_{12}k_3 + m_1 \tag{11.65}$$

$$h_2 = c_{21}k_1 + c_{22}k_3 + m_2. \tag{11.66}$$

The matrix of coefficients must be an invertible 2×2 matrix, since otherwise we could again find a nontrivial solution of (11.64). Consequently, we can solve

$$k_1 = c'_{11}h_1 + c'_{12}h_2 + m'_1 \tag{11.67}$$

$$k_3 = c'_{21}h_1 + c'_{22}h_2 + m'_2. \tag{11.68}$$

which, together with (11.62), proves the vector space decomposition

$$L^2 \cap C^{4,\alpha}_{\delta} = Ker(H') \oplus \mathcal{D}. \tag{11.69}$$

Clearly, this proves that $H' : \mathbb{R} \times \mathbb{R} \times \mathcal{D} \to C^{0,\alpha}_{\delta-4}$ is also surjective.

Finally, the estimate on the size of θ follows from Proposition 10.6:

$$\begin{aligned} \|H(0,0,0)\|_{C^{0,\alpha}_{\delta-4}} &= \|P(0)\|_{C^{0,\alpha}_{\delta-4}} = \|B^t(g^{(1)}_{a,b})\|_{C^{0,\alpha}_{\delta-4}} \\ &\leq \lambda a^4 \|k_1\|_{C^{0,\alpha}_{\delta-4}} + Ca^{4+\delta-\epsilon} \leq Ca^{4+\delta-\epsilon}. \end{aligned} \tag{11.70}$$

□

12. Completion of proofs

The following result immediately implies Theorem 1.1:

Theorem 12.1. *Let $a = b$ and $\theta \in \mathcal{D}$ be the unique solution of (11.47):*

$$P_{g^{(1)}}(\theta) = \lambda_1 k_1 + \lambda_2 k_2.$$

Then

$$\lambda_1 = \lambda a^4 + O(a^{6-\epsilon}) \tag{12.1}$$

as $a \to 0$, where

$$\lambda = \Big(\frac{2}{3} W(y_0) \circledast W(z_0) + 4tR(z_0)\mathrm{mass}(g_N)\Big)\omega_3. \tag{12.2}$$

Proof. Let $\theta \in \mathcal{D}$ be a solution of (11.47):

$$P_{g^{(1)}}(\theta) = \lambda_1 k_1 + \lambda_2 k_2.$$

Pairing both sides with ηo_1, where η is given in (11.7), and integrating (all with respect to the metric $g = g^{(1)}_a$) gives

$$\int \langle P(\theta), \eta o_1 \rangle dV = \lambda_1 \int \langle k_1, \eta o_1 \rangle dV + \lambda_2 \int \langle k_2, \eta o_1 \rangle dV. \tag{12.3}$$

The last integral is identically zero by (10.57), and by (10.43), we obtain

$$\lambda_1 = \int \langle P(\theta), \eta o_1 \rangle dV. \tag{12.4}$$

Using Proposition 3.7 we expand $P(\theta)$ as

$$\begin{aligned} P(\theta) &= P(0) + S(\theta) + Q(\theta) \\ &= B^t(g^{(1)}) + S(\theta) + Q(\theta). \end{aligned}$$

Substituting this into (12.4),

$$\lambda_1 = \int \langle B^t(g^{(1)}), \eta o_1 \rangle dV + \int \langle S(\theta), \eta o_1 \rangle dV + \int \langle Q(\theta), \eta o_1 \rangle dV. \tag{12.5}$$

Using (11.24) (replacing $\tilde{h}_i$ with θ in that computation), we estimate

$$\int \langle S(\theta), \eta o_1 \rangle dV = O(a^{2-\delta}) O(a^{4+\delta-\epsilon}) = O(a^{6-\epsilon}) \tag{12.6}$$

as $a \to 0$. The estimate (11.46) implies the pointwise estimates:

$$|\nabla^m \theta| \leq C a^{4+\delta-\epsilon} w^{\delta-m}, \tag{12.7}$$

for $0 \leq m \leq 4$. Using Proposition 3.7, the nonlinear term in (12.5) is then estimated

$$\int \langle Q(\theta), \eta o_1 \rangle dV = O(a^{8+2\delta-2\epsilon}). \tag{12.8}$$

We conclude

$$\lambda_1 = \int \langle B^t(g^{(1)}), \eta o_1 \rangle dV + O(a^{6-\epsilon}), \tag{12.9}$$

as $a \to 0$.

Notice that from (10.12),

$$\int_{DZ} \langle B^t(g^{(1)}), \eta o_1 \rangle dV = O(a^{6-\epsilon}), \tag{12.10}$$

so we can rewrite (12.9) as

$$\lambda_1 = \int_B \langle B^t(g^{(1)}), \eta o_1 \rangle dV + O(a^{6-\epsilon}), \tag{12.11}$$

as $a \to 0$, where B is the same as in (8.6).

On $B \subset N$, $\eta \equiv 1$, and from Proposition 10.4 we have

$$\begin{aligned} \int_B \langle B^t(g^{(1)}), o_1 \rangle \, dV &= \int_B \langle a^4 \lambda k_1^{(0)} - a^4 \mathcal{K}_{g_N} \delta_{g_N} \mathcal{K}_{g_N} \delta_{g_N} \overset{\circ}{\tilde{H}}_2 + O(a^8), o_1 \rangle dV \\ &= a^4 \lambda \int_B \langle k_1^{(0)}, o_1 \rangle \, dV - a^4 \int_B \langle \mathcal{K}_{g_N} \delta_{g_N} \mathcal{K}_{g_N} \delta_{g_N} \overset{\circ}{\tilde{H}}_2, o_1 \rangle \, dV \\ &\quad + \int_B \langle O(a^8), o_1 \rangle \, dV. \end{aligned} \tag{12.12}$$

By Claim 10.8, the first integral in (12.12) is

$$\begin{aligned} a^4\lambda \int_B \langle k_1^{(0)}, o_1\rangle \; dV &= a^4\lambda \int_B \langle \big(1+O(a^2)\big)k_1, o_1\rangle dV \\ &= a^4\lambda \int \langle k_1, o_1\rangle \; dV + O(a^6) \\ &= a^4\lambda + O(a^6). \end{aligned} \tag{12.13}$$

To estimate the second integral in (12.12), we use the fact that on $B \subset N$,

$$g = g_N + O(a^2).$$

In particular, for tensors T_1, T_2 we have

$$\begin{aligned} \langle T_1, T_2\rangle &= \big(1+O(a^2)\big)\langle T_1, T_2\rangle_{g_N}, \\ dV &= \big(1+O(a^2)\big)dV_{g_N}. \end{aligned}$$

Therefore,

$$\begin{aligned} -a^4 \int_B & \langle \mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2, o_1\rangle \; dV \\ &= -a^4 \int_B \langle \mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2, o_1\rangle_{g_N} \; dV_{g_N} \\ &\quad + O(a^6)\int_B \big|\mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2\big|_{g_N}\big|o_1\big|_{g_N} \; dV_{g_N}. \end{aligned} \tag{12.14}$$

For the second integral above on the right-hand side of (12.14), we note that

$$\big|\mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2\big|_{g_N} = O(|x|^{\epsilon-4}),$$

see the proof of Proposition 10.4. Also, by Theorem 6.1, o_1 decays quadratically, hence

$$\int_B \big|\mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2\big|_{g_N}\big|o_1\big|_{g_N} \; dV_{g_N} = O(1).$$

For the first integral on the right in (12.14) we recall from Theorems 5.1 and 6.1 that the trace-free part of o_1 is given by

$$\begin{aligned} \mathcal{K}_{g_N}\omega_1 &= \frac{2}{3}W_{ikj\ell}(y_0)\frac{x^k x^\ell}{|x|^4} + O(|x|^{-4+\epsilon}) \\ &= O(|x|^{-2}) \end{aligned} \tag{12.15}$$

as $|x| \to \infty$. Moreover,

$$\Box_{g_N}\omega_1 = \delta_{g_N}\mathcal{K}_{g_N}\omega_1 = 0. \tag{12.16}$$

Therefore, integration by parts gives

$$\begin{aligned}
-a^4 \int_B &\langle \mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2, o_1\rangle_{g_N}\ dV_{g_N} \\
&= -a^4 \int_B \langle \mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2, \mathcal{K}_{g_N}\omega_1\rangle_{g_N}\ dV_{g_N} \\
&= 2a^4 \int_B \langle \delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2, \delta_{g_N}\mathcal{K}_{g_N}\omega_1\rangle_{g_N}\ dV_{g_N} \\
&\quad - 2a^4 \oint_{\partial B} \mathcal{K}_{g_N}\omega_1\big(N, \delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2\big)\ dS \\
&= -2a^4 \oint_{\partial B} \mathcal{K}_{g_N}\omega_1\big(N, \delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2\big)\ dS.
\end{aligned} \tag{12.17}$$

Using (12.15), the integrand of the boundary integral above is

$$\big|\mathcal{K}_{g_N}\omega_1\big(N, \delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2\big)\big| = O(|x|^{-2}) \cdot O(|x|^{\epsilon-3}) = O(|x|^{\epsilon-5}),$$

and it follows that the boundary integral in (12.17) is of the order

$$-2a^4 \oint_{\partial B} \mathcal{K}_{g_N}\omega_1\big(N, \delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2\big)\ dS = O(a^{6-\epsilon}).$$

Consequently,

$$-a^4 \int_B \langle \mathcal{K}_{g_N}\delta_{g_N}\mathcal{K}_{g_N}\delta_{g_N}\overset{\circ}{\tilde{H}}_2, o_1\rangle\ dV = O(a^{6-\epsilon}). \tag{12.18}$$

We can also use the fact that o_1 decays quadratically to estimate the last term in (12.12) as

$$\int_B \langle O(a^8), o_1\rangle\ dV = O(a^6). \tag{12.19}$$

Combining (12.12), (12.13), (12.18) , and (12.19) we obtain

$$\int_B \langle B^t(g^{(1)}), \eta o_1\rangle dV = \lambda a^4 + O(a^{6-\epsilon}). \tag{12.20}$$

Proposition 8.1 then completes the proof. □

Proof of Theorem 1.2. From Theorem 12.1, it is clear that for a sufficiently small, there are two possibilities. The first is that the remainder term in (12.1) is identically zero for all a sufficiently small. Choosing t_0 as in (1.20), we have that $\lambda_1 = 0$. The second possibility is that the remainder term in (12.1) is not zero. In this case, by an application of the intermediate value theorem, we may perturb t slightly to again conclude that $\lambda_1 = 0$. We now have a solution of the equation

$$P_{g^{(1)}}(\theta) = \lambda_2 k_2. \tag{12.21}$$

Recalling the definition of P, this is

$$B^t(g^{(1)}+\theta)+\mathcal{K}_{g^{(1)}+\theta}\delta_{g^{(1)}}\mathcal{K}_{g^{(1)}}\delta_{g^{(1)}}\mathring{\theta}=\lambda_2 k_2. \tag{12.22}$$

With respect to the metric $g^{(1)}+\theta$, the trace of the left hand side of this equation has mean value zero, so we have

$$0=\lambda_2\int tr_{g^{(1)}+\theta}k_2 dV_{g^{(1)}+\theta}. \tag{12.23}$$

Since k_2 has compact support in the region where the weight function is bounded, expanding the trace and volume element and using (12.7), we have

$$0=\lambda_2\Big(\int tr_{g^{(1)}}k_2 dV_{g^{(1)}}+O(a^{4+\delta-\epsilon})\Big)\Big)=\lambda_2(1+O(a^{4+\delta-\epsilon})), \tag{12.24}$$

as $a\to 0$, by (10.56), which implies that $\lambda_2=0$. We have therefore found a solution of

$$P_{g^{(1)}}(\theta)=0, \tag{12.25}$$

which is a smooth B^t-flat metric from Proposition 3.5.

In the cases of multiple gluing points, imposing the bilateral, trilateral, or quadrilateral symmetries in the respective cases, reduces the argument to that of a single gluing point, so the argument is the same as above. $\square$

Proof of Theorem 1.4. In the Bach-flat case, we may restrict all above arguments to pointwise traceless tensors. The pure-trace kernel and cokernel elements are then not required in the Lyapunov-Schmidt reduction in Section 11. We then add a 1-dimensional kernel parameter to the map H. That is, we let

$$\mathcal{D}=\Big\{h\in C^{4,\alpha}_{\delta}\ :\ \int\langle h,k_1\rangle=0\Big\}. \tag{12.26}$$

where k_1 is of compact support chosen to pair non-trivially with $\mathring{o}_1$, and define the mapping $H:\mathbb{R}\times\mathbb{R}\times\mathcal{D}\to C^{\alpha}_{\delta-4}$ by

$$H(s,\lambda_1,\theta)=P_{g^{(1)}}(\theta+s\eta\mathring{o}_1)-\lambda_1 k_1. \tag{12.27}$$

For gluing parameter a sufficiently small, the Kuranishi map is then the map

$$\Psi: s\mapsto\lambda_1(s) \tag{12.28}$$

Using the gauging argument from [GV11, Section 2.3], the fixed point argument in Section 11 is easily extended to show that *any* equivariant Bach-flat metric in a sufficiently small $C^{4,\alpha}$-neighborhood of the approximate metric will correspond to a zero of Ψ for some s. If $W(y_0)\circledast W(z_0)\neq 0$, then the leading term of Ψ is non-zero, so obviously there can be no equivariant Bach-flat metric in a sufficiently small $C^{4,\alpha}$ neighborhood of the approximate metric. $\square$

12.1. **Computation of values in Table 1.1.** Assume $W^{\pm}, \widetilde{W}^{\pm}$ are trace-free endomorphisms of $\Lambda^{\pm}_2(V^*)$, where V is a real, oriented, four-dimensional inner product space, and write

$$\begin{aligned} W &= W^+ + W^- : \Lambda^2(V^*) \to \Lambda^2(V^*), \\ \widetilde{W} &= \widetilde{W}^+ + \widetilde{W}^- : \Lambda^2(V^*) \to \Lambda^2(V^*). \end{aligned}$$

Assume further that W and $\widetilde{W}$ can be simultaneously diagonalized: that is, there is an orthogonal basis of eigenvectors (two-forms) for W and $\widetilde{W}$ denoted

$$\omega, \eta, \theta, \omega^-, \eta^-, \theta^-, \tag{12.29}$$

where the first three are a basis of $\Lambda^+_2(V^*)$ and the last three a basis of $\Lambda^-_2(V^*)$. Denote the eigenvalues of W and $\widetilde{W}$ as

$$\begin{aligned} \mathrm{spec}(W) &= \{\lambda, \mu, \nu, \lambda^-, \mu^-, \nu^-\}, \\ \mathrm{spec}(\widetilde{W}) &= \{\tilde{\lambda}, \tilde{\mu}, \tilde{\nu}, \tilde{\lambda}^-, \tilde{\mu}^-, \tilde{\nu}^-\}. \end{aligned} \tag{12.30}$$

We will further assume that $W^{\pm}$ and $\widetilde{W}^{\pm}$ are trace-free; i.e.,

$$\begin{aligned} &\lambda + \mu + \nu = 0,\ \lambda^- + \mu^- + \nu^- = 0, \\ &\tilde{\lambda} + \tilde{\mu} + \tilde{\nu} = 0,\ \tilde{\lambda}^- + \tilde{\mu}^- + \tilde{\nu}^- = 0. \end{aligned} \tag{12.31}$$

Using this basis in (12.29) we can write

$$\begin{aligned} W = \frac{1}{2}\big\{ &\lambda\omega \otimes \omega + \mu\eta \otimes \eta + \nu\theta \otimes \theta \\ &+ \lambda^-\omega^- \otimes \omega^- + \mu^-\eta^- \otimes \eta^- + \nu^-\theta^- \otimes \theta^-\big\}, \\ \widetilde{W} = \frac{1}{2}\big\{ &\tilde{\lambda}\omega \otimes \omega + \tilde{\mu}\eta \otimes \eta + \tilde{\nu}\theta \otimes \theta \\ &+ \tilde{\lambda}^-\omega^- \otimes \omega^- + \tilde{\mu}^-\eta^- \otimes \eta^- + \tilde{\nu}^-\theta^- \otimes \theta^-\big\}. \end{aligned} \tag{12.32}$$

We normalize the eigenforms to have length $\sqrt{2}$; this convention gives the identities

$$\omega^2 = \eta^2 = \cdots = (\theta^-)^2 = Id. \tag{12.33}$$

We also point out two more important algebraic facts: first, the product of any SD basis element with any ASD basis element gives a symmetric trace-free two-tensor, whose square is the identity. Thus, for example,

$$h = \omega\omega^- \Rightarrow\ tr\ h = 0,\ h^2 = Id,\ |h|^2 = 4. \tag{12.34}$$

Also, the bases of $\Lambda^{\pm}_2$ give a quaternionic structure satisfying the following multiplication rules:

$$\omega\eta = \theta,\ \eta\theta = \omega,\ \theta\omega = \eta. \tag{12.35}$$

Lemma 12.2. *Fix an orthonormal basis* $\{v_1, \dots, v_4\}$ *of* V*, and let* $W_{ikj\ell}$ *(resp.,* $\widetilde{W}_{i\ell jk}$*) denote the components of* W *(resp.,* $\widetilde{W}$*) with respect to this basis. Then*

$$\begin{aligned} W_{ikj\ell}\widetilde{W}_{i\ell jk} &= 2\big\{\lambda\tilde{\lambda} + \mu\tilde{\mu} + \nu\tilde{\nu} + \lambda^-\tilde{\lambda}^- + \mu^-\tilde{\mu}^- + \nu^-\tilde{\nu}^-\big\} \\ &= 2\langle W, \widetilde{W}\rangle = \frac{1}{2}W_{ikj\ell}\widetilde{W}_{ikj\ell}. \end{aligned} \tag{12.36}$$

In particular, the answer is independent of the choice of basis.

Proof. By (12.32),

$$\begin{aligned} 4W_{ikj\ell}\widetilde{W}_{i\ell jk} = \big\{&\lambda\omega_{ik}\omega_{j\ell} + \mu\eta_{ik}\eta_{j\ell} + \nu\theta_{ik}\theta_{j\ell} \\ &+ \lambda^-\omega^-_{ik}\omega^-_{j\ell} + \mu^-\eta^-_{ik}\eta^-_{j\ell} + \nu^-\theta^-_{ik}\theta^-_{j\ell}\big\} \\ \times\big\{&\tilde{\lambda}\omega_{i\ell}\omega_{jk} + \tilde{\mu}\eta_{i\ell}\eta_{jk} + \tilde{\nu}\theta_{i\ell}\theta_{jk} \\ &+ \tilde{\lambda}^-\omega^-_{i\ell}\omega^-_{jk} + \tilde{\mu}^-\eta^-_{i\ell}\eta^-_{jk} + \tilde{\nu}^-\theta^-_{i\ell}\theta^-_{jk}\big\}. \end{aligned} \tag{12.37}$$

As we multiply and distribute we see that there are six kinds of terms, which we represent schematically as

$$\begin{aligned} &(A_{ik}A_{j\ell})\cdot(A_{i\ell}A_{jk}), \ \ (A^-_{ik}A^-_{j\ell}\cdot(A^-_{i\ell}A^-_{jk}) \\ &(A_{ik}A_{j\ell})\cdot(B_{i\ell}B_{jk}), \ \ (A^-_{ik}A^-_{j\ell})\cdot(B^-_{i\ell}B^-_{jk}) \\ &(A_{ik}A_{j\ell})\cdot(C^-_{i\ell}C^-_{jk}), \ \ (A^-_{ik}A^-_{j\ell})\cdot(C_{i\ell}C_{jk}), \end{aligned} \tag{12.38}$$

where A, B, C are self-dual and A^-, B^-, and C^- are anti-self-dual. Using the multiplication rules in (12.33) and (12.34), we find

$$\begin{aligned} (A_{ik}A_{j\ell})\cdot(A_{i\ell}A_{jk}) = 4, &\ \ (A^-_{ik}A^-_{j\ell}\cdot(A^-_{i\ell}A^-_{jk}) = 4, \\ (A_{ik}A_{j\ell})\cdot(B_{i\ell}B_{jk}) = -4, &\ \ (A^-_{ik}A^-_{j\ell})\cdot(B^-_{i\ell}B^-_{jk}) = -4, \\ (A_{ik}A_{j\ell})\cdot(C^-_{i\ell}C^-_{jk}) = -4, &\ \ (A^-_{ik}A^-_{j\ell})\cdot(C_{i\ell}C_{jk}) = -4. \end{aligned} \tag{12.39}$$

Therefore, after multiplying out and collecting all the terms in (12.37), we find

$$\begin{aligned} 4W_{ikj\ell}\widetilde{W}_{i\ell jk} &= 4\lambda(\tilde{\lambda} - \tilde{\mu} - \tilde{\nu}) + 4\lambda(-\tilde{\lambda}^- - \tilde{\mu}^- - \tilde{\nu}^-) \\ &+ 4\mu(-\tilde{\lambda} + \tilde{\mu} - \tilde{\nu}) + 4\mu(-\tilde{\lambda}^- - \tilde{\mu}^- - \tilde{\nu}^-) \\ &+ 4\nu(-\tilde{\lambda} - \tilde{\mu} + \tilde{\nu}) + 4\nu(-\tilde{\lambda}^- - \tilde{\mu}^- - \tilde{\nu}^-) \\ &+ 4\lambda^-(-\tilde{\lambda} - \tilde{\mu} - \tilde{\nu}) + 4\lambda^-(\tilde{\lambda}^- - \tilde{\mu}^- - \tilde{\nu}^-) \\ &+ 4\mu^-(-\tilde{\lambda} - \tilde{\mu} - \tilde{\nu}) + 4\mu^-(-\lambda^- + \mu^- - \nu^-) \\ &+ 4\nu^-(-\tilde{\lambda} - \tilde{\mu} - \tilde{\nu}) + 4\nu^-(-\tilde{\lambda}^- - \tilde{\mu}^- + \tilde{\nu}^-). \end{aligned} \tag{12.40}$$

By (12.31), this gives

$$\begin{aligned} 4W_{ikj\ell}\widetilde{W}_{i\ell jk} &= 8\big\{\lambda\tilde{\lambda} + \mu\tilde{\mu} + \nu\tilde{\nu} + \lambda^-\tilde{\lambda}^- + \mu^-\tilde{\mu}^- + \nu^-\tilde{\nu}^-\big\} \\ &= 8\langle W, \widetilde{W}\rangle, \end{aligned} \tag{12.41}$$

and (12.36) follows. □

To compute the values of t_0, we note that in the coordinate system $\{z^i\}$ given in Section 2, letting

$$\begin{aligned}\omega^{\pm} &= e^1 \wedge e^2 \pm e^3 \wedge e^4,\\ \eta^{\pm} &= e^1 \wedge e^3 \mp e^2 \wedge e^4,\\ \theta^{\pm} &= e^1 \wedge e^4 \pm e^2 \wedge e^3,\end{aligned}$$

with $(e^1, e^3, e^3, e^4) = (dz^1, dz^2, dz^3, dz^4)$, we have

$$\begin{aligned}W^+(g_{FS}) &= \mathrm{diag}(R/6, -R/12, -R/12) = \mathrm{diag}(4, -2, -2),\\ W^-(g_{FS}) &= \mathrm{diag}(0, 0, 0),\\ W^+(g_{S^2\times S^2}) &= \mathrm{diag}(R/6, -R/12, -R/12) = \mathrm{diag}(2/3, -1/3, -1/3),\\ W^-(g_{S^2\times S^2}) &= \mathrm{diag}(2/3, -1/3, -1/3).\end{aligned}$$

In case (i), since $\mathrm{mass}(\hat{g}_{FS}) = 2$,

$$t_0 = \frac{-1}{6\cdot 24\cdot 2}4(16+4+4) = -\frac{1}{3}. \tag{12.42}$$

In case (ii) with a Burns metric attached, we have

$$t_0 = \frac{-1}{6\cdot 4\cdot 2}4(4\cdot(2/3) + 2\cdot(1/3) + 2\cdot(1/3)) = -\frac{1}{3}. \tag{12.43}$$

Case (v) has the same value as this.

In case (ii) with a Green's function $S^2 \times S^2$ attached,

$$\begin{aligned}t_0 &= \frac{-1}{6\cdot 24\cdot \mathrm{mass}(\hat{g}_{S^2\times S^2})}4((2/3)\cdot 4 + (1/3)\cdot 2 + (1/3)\cdot 2)\\ &= -\frac{1}{9\cdot \mathrm{mass}(\hat{g}_{S^2\times S^2})}.\end{aligned} \tag{12.44}$$

In case (iii),

(12.45)

$$\begin{aligned}t_0 &= \frac{-1}{6\cdot 4\cdot \mathrm{mass}(\hat{g}_{S^2\times S^2})}((2/3)^2 + (1/3)^2 + (1/3)^2 + (2/3)^2 + (1/3)^2 + (1/3)^2)\\ &= -\frac{2}{9\cdot \mathrm{mass}(\hat{g}_{S^2\times S^2})}.\end{aligned}$$

Case (iv) has the same value of t_0 as does case (iii).

All other cases are computed similarly as the above cases, so it is not necessary to write every case here. We only need mention the fact that in all non-orientable cases, the answer does not depend on choice of local orientation.

Appendix A. Proof of Proposition 8.5

To prove the proposition, we use the expansion of the metric in AF coordinates,

$$g_{\mu\nu} = \delta_{\mu\nu} - \frac{1}{3}R_{\mu k\nu \ell}(y_0)\frac{x^k x^\ell}{|x|^4} + \frac{2A}{|x|^2}\delta_{\mu\nu} + O(|x|^{-3}). \tag{A.1}$$

In terms of the Christoffel symbols, the Ricci tensor is given by

$$\begin{aligned} R_{ij} &= \partial_m \Gamma^m_{ij} - \partial_i \Gamma^m_{jm} + \Gamma * \Gamma \\ &= -\frac{1}{2} g^{mp} \big\{ \partial_m \partial_p g_{ij} - \partial_m \partial_i g_{jp} - \partial_m \partial_j g_{ip} + \partial_i \partial_j g_{mp} \big\} + \partial g * \partial g. \end{aligned} \tag{A.2}$$

By the expansion (A.1),

$$g^{mp} = \delta_{mp} + O(|x|^{-2}), \tag{A.3}$$

hence

$$\begin{aligned} R_{ij} &= -\frac{1}{2} \big\{ \delta_{mp} + O(|x|^{-2}) \big\} \big\{ \partial_m \partial_p g_{ij} - \partial_m \partial_i g_{jp} - \partial_m \partial_j g_{ip} + \partial_i \partial_j g_{mp} \big\} + \partial g * \partial g \\ &= -\frac{1}{2} \big\{ \partial_m \partial_m g_{ij} - \partial_m \partial_i g_{jm} - \partial_m \partial_j g_{im} + \partial_i \partial_j g_{mm} \big\} + (\partial^2 g) * (|x|^{-2}) + \partial g * \partial g. \end{aligned} \tag{A.4}$$

In AF coordinates, $\partial g = O(|x|^{-3}), \partial^2 g = O(|x|^{-4})$; hence

$$R_{ij} = -\frac{1}{2} \big\{ \partial_m \partial_m g_{ij} - \partial_m \partial_i g_{jm} - \partial_m \partial_j g_{im} + \partial_i \partial_j g_{mm} \big\} + O(|x|^{-6}). \tag{A.5}$$

By (A.1),

$$\begin{aligned} \partial_\beta g_{\mu\nu} &= -\frac{1}{3} R_{\mu\beta\nu k}(y_0) \frac{x^k}{|x|^4} - \frac{1}{3} R_{\mu k \nu \beta}(y_0) \frac{x^k}{|x|^4} \\ &\quad + \frac{4}{3} R_{\mu k \nu \ell}(y_0) \frac{x^k x^\ell x^\beta}{|x|^6} - \frac{4A}{|x|^4} x^\beta \delta_{\mu\nu} + O(|x|^{-4}), \end{aligned} \tag{A.6}$$

$$\begin{aligned} \partial_\alpha \partial_\beta g_{\mu\nu} &= -\frac{1}{3} R_{\mu\beta\nu\alpha}(y_0) \frac{1}{|x|^4} - \frac{1}{3} R_{\mu\alpha\nu\beta}(y_0) \frac{1}{|x|^4} \\ &\quad + \frac{4}{3} R_{\mu\beta\nu k}(y_0) \frac{x^k x^\alpha}{|x|^6} + \frac{4}{3} R_{\mu k \nu \beta}(y_0) \frac{x^k x^\alpha}{|x|^6} \\ &\quad + \frac{4}{3} R_{\mu\alpha\nu k}(y_0) \frac{x^k x^\beta}{|x|^6} + \frac{4}{3} R_{\mu k \nu \alpha}(y_0) \frac{x^k x^\beta}{|x|^6} \\ &\quad + \frac{4}{3} R_{\mu k \nu \ell}(y_0) \frac{x^k x^\ell}{|x|^6} \delta_{\alpha\beta} - 8 R_{\mu k \nu \ell}(y_0) \frac{x^k x^\ell x^\alpha x^\beta}{|x|^8} \\ &\quad - \frac{4A}{|x|^4} \delta_{\alpha\beta} \delta_{\mu\nu} + \frac{16A}{|x|^6} x^\alpha x^\beta \delta_{\mu\nu} + O(|x|^{-5}). \end{aligned} \tag{A.7}$$

Consequently, the first term in (A.5) is

$$\begin{aligned}\partial_m\partial_m g_{ij} &= -\frac{1}{3}R_{imjm}(y_0)\frac{1}{|x|^4} - \frac{1}{3}R_{imjm}(y_0)\frac{1}{|x|^4}\\ &\quad + \frac{4}{3}R_{imjk}(y_0)\frac{x^kx^m}{|x|^6} + \frac{4}{3}R_{ikjm}(y_0)\frac{x^kx^m}{|x|^6}\\ &\quad + \frac{4}{3}R_{imjk}(y_0)\frac{x^kx^m}{|x|^6} + \frac{4}{3}R_{ikjm}(y_0)\frac{x^kx^m}{|x|^6}\\ &\quad + \frac{4}{3}R_{ikj\ell}(y_0)\frac{x^kx^\ell}{|x|^6}\delta_{mm} - 8R_{ikj\ell}(y_0)\frac{x^kx^\ell x^mx^m}{|x|^8}\\ &\quad - \frac{4A}{|x|^4}\delta_{mm}\delta_{ij} + \frac{16A}{|x|^6}x^mx^m\delta_{ij} + O(|x|^{-5}).\end{aligned}\tag{A.8}$$

The first two terms combine to give a Ricci curvature term, while the third through the eighth terms are all the same (though with different coefficients); adding up we get

$$\partial_m\partial_m g_{ij} = -\frac{2}{3}R_{ij}(y_0)\frac{1}{|x|^4} + \frac{8}{3}R_{ikj\ell}(y_0)\frac{x^kx^\ell}{|x|^6} + O(|x|^{-5}).\tag{A.9}$$

The second term is

$$\begin{aligned}\partial_m\partial_i g_{jm} &= -\frac{1}{3}R_{jimm}(y_0)\frac{1}{|x|^4} - \frac{1}{3}R_{jmmi}(y_0)\frac{1}{|x|^4}\\ &\quad + \frac{4}{3}R_{jimk}(y_0)\frac{x^kx^m}{|x|^6} + \frac{4}{3}R_{jkmi}(y_0)\frac{x^kx^m}{|x|^6}\\ &\quad + \frac{4}{3}R_{jmmk}(y_0)\frac{x^kx^i}{|x|^6} + \frac{4}{3}R_{jkmm}(y_0)\frac{x^kx^i}{|x|^6}\\ &\quad + \frac{4}{3}R_{jkm\ell}(y_0)\frac{x^kx^\ell}{|x|^6}\delta_{mi} - 8R_{jkm\ell}(y_0)\frac{x^kx^\ell x^mx^i}{|x|^8}\\ &\quad - \frac{4A}{|x|^4}\delta_{mi}\delta_{jm} + \frac{16A}{|x|^6}x^mx^i\delta_{jm} + O(|x|^{-5}),\end{aligned}\tag{A.10}$$

notice that the fourth and seventh terms cancel each other, while the first, sixth, and eight terms vanish because of the skew-symmetry of the curvature tensor. Also, the second and fifth terms are traces. Therefore,

$$\partial_m\partial_i g_{jm} = \frac{1}{3}R_{ij}(y_0)\frac{1}{|x|^4} - \frac{4}{3}R_{jk}(y_0)\frac{x^ix^k}{|x|^6} - \frac{4A}{|x|^4}\delta_{ij} + \frac{16A}{|x|^6}x^ix^j + O(|x|^{-5}),\tag{A.11}$$

while the third term is

$$\partial_m\partial_j g_{im} = \frac{1}{3}R_{ij}(y_0)\frac{1}{|x|^4} - \frac{4}{3}R_{ik}(y_0)\frac{x^jx^k}{|x|^6} - \frac{4A}{|x|^4}\delta_{ij} + \frac{16A}{|x|^6}x^ix^j + O(|x|^{-5}).\tag{A.12}$$

The last term in (A.5) is

$$\begin{aligned}\partial_i\partial_j g_{mm} = &-\frac{1}{3}R_{mjmi}(y_0)\frac{1}{|x|^4} - \frac{1}{3}R_{mimj}(y_0)\frac{1}{|x|^4}\\ &+\frac{4}{3}R_{mjmk}(y_0)\frac{x^kx^i}{|x|^6} + \frac{4}{3}R_{mkmj}(y_0)\frac{x^kx^i}{|x|^6}\\ &+\frac{4}{3}R_{mimk}(y_0)\frac{x^kx^j}{|x|^6} + \frac{4}{3}R_{mkmi}(y_0)\frac{x^kx^j}{|x|^6}\\ &+\frac{4}{3}R_{mkm\ell}(y_0)\frac{x^kx^\ell}{|x|^6}\delta_{ij} - 8R_{mkm\ell}(y_0)\frac{x^kx^\ell x^ix^j}{|x|^8}\\ &-\frac{4A}{|x|^4}\delta_{ij}\delta_{mm} + \frac{16A}{|x|^6}x^ix^j\delta_{mm} + O(|x|^{-5}).\end{aligned}\tag{A.13}$$

In this case all the curvature terms involve traces, so we get

$$\begin{aligned}\partial_i\partial_j g_{mm} = &-\frac{2}{3}R_{ij}(y_0)\frac{1}{|x|^4} + \frac{8}{3}R_{jk}(y_0)\frac{x^ix^k}{|x|^6} + \frac{8}{3}R_{ik}(y_0)\frac{x^jx^k}{|x|^6}\\ &+\frac{4}{3}R_{k\ell}(y_0)\frac{x^kx^\ell}{|x|^6}\delta_{ij} - 8R_{k\ell}(y_0)\frac{x^kx^\ell x^ix^j}{|x|^8}\\ &-\frac{16A}{|x|^4}\delta_{ij} + \frac{64A}{|x|^4}x^ix^j + O(|x|^{-5}).\end{aligned}\tag{A.14}$$

Combining (A.9)–(A.14),

$$\begin{aligned}&\partial_m\partial_m g_{ij} - \partial_m\partial_i g_{jm} - \partial_m\partial_j g_{im} + \partial_i\partial_j g_{mm} =\\ &\qquad\frac{8}{3}R_{ikj\ell}(y_0)\frac{x^kx^\ell}{|x|^6} - 2R_{ij}(y_0)\frac{1}{|x|^4} + 4R_{jk}(y_0)\frac{x^ix^k}{|x|^6} + 4R_{ik}(y_0)\frac{x^jx^k}{|x|^6}\\ &\qquad+\frac{4}{3}R_{k\ell}(y_0)\frac{x^kx^\ell}{|x|^6}\delta_{ij} - 8R_{k\ell}(y_0)\frac{x^kx^\ell x^ix^j}{|x|^8} - \frac{8A}{|x|^4}\delta_{ij} + \frac{32A}{|x|^6}x^ix^j + O(|x|^{-5}).\end{aligned}\tag{A.15}$$

We now use the fact that (Y, g_Y) is Einstein, and that $\{y^i\}$ are normal coordinates centered at y_0:

$$\begin{aligned}R_{ij}(y_0) &= \frac{1}{4}R(y_0)\delta_{ij},\\ R_{ijk\ell}(y_0) &= W_{ijk\ell}(y_0) + \frac{1}{12}R(y_0)\big(\delta_{ik}\delta_{j\ell} - \delta_{i\ell}\delta_{jk}\big).\end{aligned}$$

Substituting these gives

$$\begin{aligned}&\partial_m\partial_m g_{ij} - \partial_m\partial_i g_{jm} - \partial_m\partial_j g_{im} + \partial_i\partial_j g_{mm} =\\ &\qquad\frac{8}{3}W_{ikj\ell}(y_0)\frac{x^kx^\ell}{|x|^6} + \frac{1}{18}R(y_0)\frac{1}{|x|^4}\delta_{ij} - \frac{2}{9}R_{(}y_0)\frac{x^ix^j}{|x|^6}\\ &\qquad-\frac{8A}{|x|^4}\delta_{ij} + \frac{32A}{|x|^6}x^ix^j + O(|x|^{-5}),\end{aligned}\tag{A.16}$$

and (8.43) follows.

Appendix B. Non-simply-connected examples

Table B.1. Non-simply-connected examples with more than one bubble

Topology of connected sum	Value of t_0	Symmetry
$S^2 \times S^2 \# 2(S^2 \times S^2/\mathbb{Z}_2)$	$-2(9m_2)^{-1}$	bilateral
$S^2 \times S^2 \# 2(\mathbb{RP}^2 \times \mathbb{RP}^2)$	$-2(9m_3)^{-1}$	bilateral
$\mathbb{CP}^2 \# 3(S^2 \times S^2/\mathbb{Z}_2)$	$-(9m_2)^{-1}$	trilateral
$\mathbb{CP}^2 \# 3(\mathbb{RP}^2 \times \mathbb{RP}^2)$	$-(9m_3)^{-1}$	trilateral
$S^2 \times S^2 \# 4(S^2 \times S^2/\mathbb{Z}_2)$	$-2(9m_2)^{-1}$	quadrilateral
$S^2 \times S^2 \# 4(\mathbb{RP}^2 \times \mathbb{RP}^2)$	$-2(9m_3)^{-1}$	quadrilateral
$(S^2 \times S^2/\mathbb{Z}_2) \# 2\overline{\mathbb{CP}}^2$	$-1/3$	bilateral
$(S^2 \times S^2/\mathbb{Z}_2) \# 2(S^2 \times S^2)$	$-2(9m_1)^{-1}$	bilateral
$3\#(S^2 \times S^2/\mathbb{Z}_2)$	$-2(9m_2)^{-1}$	bilateral
$(S^2 \times S^2/\mathbb{Z}_2) \# 2(\mathbb{RP}^2 \times \mathbb{RP}^2)$	$-2(9m_3)^{-1}$	bilateral

All non-simply-connected possiblities with more than one bubble are listed in Table B.1. The approximate metric in each case is obtained by using the first factor as the compact manifold, with the AF space clear from the latter factors. The first two cases are analogous to Cases (iv) and (v). The third and fourth are analogous to Cases (vi) and (vii). The fifth and six are analogous to Cases (vii) and (ix). The last four cases require a short explanation. In the case of $S^2 \times S^2/\mathbb{Z}_2$, there are two fixed points: the equivalence classes of (n, n) and (n, s). The diagonal symmetry descends to the quotient, and fixes both of these points. The symmetry of reflection in a horizontal line descends to the quotient, and this interchanges the fixed points, and we again call this invariance bilateral symmetry. We may therefore glue on the same AF space at each fixed point, and require bilateral symmetry, which yields the last four cases in Table B.1.

References

[And89] Michael T. Anderson, *Ricci curvature bounds and Einstein metrics on compact manifolds*, J. Amer. Math. Soc. **2** (1989), no. 3, 455–490.

[AP06] Claudio Arezzo and Frank Pacard, *Blowing up and desingularizing constant scalar curvature Kähler manifolds*, Acta Math. **196** (2006), no. 2, 179–228.

[AP09] ———, *Blowing up Kähler manifolds with constant scalar curvature. II*, Ann. of Math. (2) **170** (2009), no. 2, 685–738.

[APS11] Claudio Arezzo, Frank Pacard, and Michael Singer, *Extremal metrics on blowups*, Duke Math. J. **157** (2011), no. 1, 1–51.

[AV12a] Antonio G. Ache and Jeff A. Viaclovsky, *Asymptotics of the self-dual deformation complex*, arXiv.org:1201.1028, 2012.

[AV12b] ———, *Obstruction-flat asymptotically locally Euclidean metrics*, Geom. Funct. Anal. **22** (2012), no. 4, 832–877.

[Bar86] Robert Bartnik, *The mass of an asymptotically flat manifold*, Comm. Pure Appl. Math. **39** (1986), no. 5, 661–693.

[Bes87] Arthur L. Besse, *Einstein manifolds*, Springer-Verlag, Berlin, 1987.

[Biq11] Olivier Biquard, *Désingularisation de métriques d'Einstein. I.*, arXiv:1105.4904, 2011.

[BKN89] Shigetoshi Bando, Atsushi Kasue, and Hiraku Nakajima, *On a construction of coordinates at infinity on manifolds with fast curvature decay and maximal volume growth*, Invent. Math. **97** (1989), no. 2, 313–349.

[Bre08] Simon Brendle, *Blow-up phenomena for the Yamabe equation*, J. Amer. Math. Soc. **21** (2008), no. 4, 951–979.

[CLW08] Xiuxiong Chen, Claude Lebrun, and Brian Weber, *On conformally Kähler, Einstein manifolds*, J. Amer. Math. Soc. **21** (2008), no. 4, 1137–1168.

[Der83] Andrzej Derdziński, *Self-dual Kähler manifolds and Einstein manifolds of dimension four*, Compositio Math. **49** (1983), no. 3, 405–433.

[DF89] S. Donaldson and R. Friedman, *Connected sums of self-dual manifolds and deformations of singular spaces*, Nonlinearity **2** (1989), no. 2, 197–239.

[EGH80] Tohru Eguchi, Peter B. Gilkey, and Andrew J. Hanson, *Gravitation, gauge theories and differential geometry*, Phys. Rep. **66** (1980), no. 6, 213–393.

[Flo91] Andreas Floer, *Self-dual conformal structures on* $l\mathbf{C}\mathrm{P}^2$, J. Differential Geom. **33** (1991), no. 2, 551–573.

[GV11] Matthew J. Gursky and Jeff A. Viaclovsky, *Rigidity and stability of Einstein metrics for quadratic curvature functionals*, arXiv.org:1105.4648, 2011.

[KN96] Shoshichi Kobayashi and Katsumi Nomizu, *Foundations of differential geometry. Vol. II*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1996, Reprint of the 1969 original, A Wiley-Interscience Publication.

[KS01] A. Kovalev and M. Singer, *Gluing theorems for complete anti-self-dual spaces*, Geom. Funct. Anal. **11** (2001), no. 6, 1229–1281.

[LeB88] Claude LeBrun, *Counter-examples to the generalized positive action conjecture*, Comm. Math. Phys. **118** (1988), no. 4, 591–596.

[LeB91] ———, *Explicit self-dual metrics on* $\mathbf{C}\mathrm{P}_2\#\cdots\#\mathbf{C}\mathrm{P}_2$, J. Differential Geom. **34** (1991), no. 1, 223–253.

[LeB12] ———, *On Einstein, Hermitian 4-manifolds*, J. Differential Geom. **90** (2012), no. 2, 277–302.

[LM85] Robert B. Lockhart and Robert C. McOwen, *Elliptic differential operators on noncompact manifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **12** (1985), no. 3, 409–447.

[LNN97] Claude LeBrun, Shin Nayatani, and Takashi Nitta, *Self-dual manifolds with positive Ricci curvature*, Math. Z. **224** (1997), no. 1, 49–63.

[LP87] John M. Lee and Thomas H. Parker, *The Yamabe problem*, Bull. Amer. Math. Soc. (N.S.) **17** (1987), no. 1, 37–91.

[LP11] H. Lü and C. N. Pope, *Critical gravity in four dimensions*, Phys. Rev. Lett. **106** (2011), 181302.

[Mal11] J. Maldacena, *Einstein gravity from conformal gravity*, arXiv:1105.5632v2, 2011.

[Pet06] Peter Petersen, *Riemannian geometry*, second ed., Graduate Texts in Mathematics, vol. 171, Springer, New York, 2006.

[Poo86] Y. Sun Poon, *Compact self-dual manifolds with positive scalar curvature*, J. Differential Geom. **24** (1986), no. 1, 97–132.

[Sch07] Hans-Jürgen Schmidt, *Fourth order gravity: equations, history, and applications to cosmology*, Int. J. Geom. Methods Mod. Phys. **4** (2007), no. 2, 209–248.

[Ste78] K. S. Stelle, *Classical gravity with higher derivatives*, General Relativity and Gravitation **9** (1978), no. 4, 353–371.

[SY79] Richard M. Schoen and Shing Tung Yau, *On the proof of the positive mass conjecture in general relativity*, Comm. Math. Phys. **65** (1979), no. 1, 45–76.

[SY81] ———, *Proof of the positive mass theorem. II*, Comm. Math. Phys. **79** (1981), no. 2, 231–260.

[Szé12] Gábor Székelyhidi, *On blowing up extremal Kähler manifolds*, Duke Math. J. **161** (2012), no. 8, 1411–1453.

[Tau92] Clifford Henry Taubes, *The existence of anti-self-dual conformal structures*, J. Differential Geom. **36** (1992), no. 1, 163–253.

[Tia90] Gang Tian, *On Calabi's conjecture for complex surfaces with positive first Chern class*, Invent. Math. **101** (1990), no. 1, 101–172.

[TV05a] Gang Tian and Jeff Viaclovsky, *Bach-flat asymptotically locally Euclidean metrics*, Invent. Math. **160** (2005), no. 2, 357–415.

[TV05b] ———, *Moduli spaces of critical Riemannian metrics in dimension four*, Adv. Math. **196** (2005), no. 2, 346–372.

[TV08] ———, *Volume growth, curvature decay, and critical metrics*, Comment. Math. Helv. **83** (2008), no. 4, 889–911.

[Via10] Jeff Viaclovsky, *Monopole metrics and the orbifold Yamabe problem*, Annales de L'Institut Fourier **60** (2010), no. 7, 2503–2543.

[Wit81] Edward Witten, *A new proof of the positive energy theorem*, Comm. Math. Phys. **80** (1981), no. 3, 381–402.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556
E-mail address: `mgursky@nd.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI 53706
E-mail address: `jeffv@math.wisc.edu`